

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 December 2001 (20.12.2001)

PCT

(10) International Publication Number  
**WO 01/96584 A2**

(51) International Patent Classification: **C12N 15/82**

(21) International Application Number: **PCT/US01/18911**

(22) International Filing Date: **12 June 2001 (12.06.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:  
**60/210,917 12 June 2000 (12.06.2000) US**

(71) Applicant (for all designated States except US):  
**AKKADIX CORPORATION [US/US]; 4204 Sorrento Valley Blvd., Suite A, San Diego, CA 92121-1412 (US).**

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MUSHEGIAN, Arcady, R. [---/US]; 3987 Santa Nella Place, San Diego, CA 92130 (US). TAYLOR, Christopher, G. [---/US]; 2910-A**

Lucimaga Street, Carlsbad, CA 92009 (US). **FEITELSON, Jerold, S. [---/US]; 4387 Mistral Place, San Diego, CA 92130 (US). EROSHKIN, Alexy, M. [---/US]; 3803 Ruelle San Rafael, San Diego, CA 92130 (US).**


(74) Agents: **LLOYD, Jeff et al.; Saliwanchik, Lloyd & Saliwanchik, Suite A-1, 2421 N.W. 41st Street, Gainesville, FL 32606 (US).**

(81) Designated States (national): **AL, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**

(84) Designated States (regional): **ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).**

[Continued on next page]

(54) Title: **MATERIALS AND METHODS FOR THE CONTROL OF NEMATODES**



(57) Abstract: The subject invention provides novel methods and compositions for controlling nematodes. More specifically, the subject invention provides RNAi molecules, polynucleotide sequences, and methods of using these sequences in nematode control.



**Published:**

- without international search report and to be republished upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## DESCRIPTION

### MATERIALS AND METHODS FOR THE CONTROL OF NEMATODES

#### Background of the Invention

[0001] Plant parasitic nematodes, such as root-knot nematodes (*Meloidogyne* species) and cyst nematodes (*Globodera* and *Heterodera*), attack nearly every food crop, and are among the world's most damaging agricultural pests. For example, root-knot nematodes parasitize more than 2,000 plant species from diverse plant families and represent a tremendous threat to crop production world-wide. These biotrophic pathogens have evolved highly specialized and complex feeding relationships with their hosts.

[0002] Nematodes cause millions of dollars of damage each year to turf grasses, ornamental plants, and food crops. Efforts to eliminate or minimize damage caused by nematodes in agricultural settings have typically involved the use of soil fumigation with materials such as chloropicrin, methyl bromide, and dazomet, which volatilize to spread the active ingredient throughout the soil. Such fumigation materials can be highly toxic and may create an environmental hazard. Various non-fumigant chemicals have also been used, but these too create serious environmental problems and can be highly toxic to humans.

[0003] Some research articles have been published concerning the effects of  $\delta$ -endotoxins from *B. thuringiensis* species on the viability of nematodes. See, for example, Bottjer, Bone and Gill ([1985] *Experimental Parasitology* 60:239-244); Ignoffo and Dropkin (Ignoffo, C.M., Dropkin, V.H. [1977] *J. Kans. Entomol. Soc.* 50:394-398); and Ciordia, H. and W.E. Bizzell ([1961] *Jour. of Parasitology* 47:41 [abstract]). Several patents have issued describing the control of nematodes with *B.t.* See, for example, U.S. Patent Nos. 4,948,734; 5,093,120; 5,281,530; 5,426,049; 5,439,881; 5,236,843; 5,322,932; 5,151,363; 5,270,448; 5,350,577; 5,667,993; and 5,670,365. The development of resistance by insects to *B.t.* toxins is one obstacle to the successful use of such toxins.

[0004] The pesticidal activity of avermectins is well known. The avermectins are disaccharide derivatives of pentacyclic, 16-membered lactones. They can be divided into four major compounds: A<sub>1a</sub>, A<sub>2a</sub>, B<sub>1a</sub>, and B<sub>2a</sub>; and four minor compounds: A<sub>1b</sub>, A<sub>2b</sub>, B<sub>1b</sub>, and B<sub>2b</sub>. The isolation and purification of these compounds is also described in U.S. Patent No. 4,310,519, issued January 12, 1982. Avermectin B<sub>2a</sub> is active against the root-knot nematode, *Meloidogyne incognita*. It is reported to be 10-30 times as potent as commercial contact nematicides when incorporated into soil at 0.16-0.25 kg/ha (Boyce Thompson Institute for Plant Research 58th Annual Report [1981]; Putter, I. *et al.* [1981] "Avermectins: Novel Insecticides, Acaracides, and Nematicides from a Soil Microorganism," *Experientia* 37:963-964). Avermectin B<sub>2a</sub> is not toxic to tomatoes or cucumbers at rates of up to 10 kg/ha.

[0005] Fatty acids are a class of natural compounds which occur abundantly in nature and which have interesting and valuable biological activities. Tarjan and Cheo (Tarjan, A.C., P.C. Cheo [1956] "Nematocidal Value of Some Fatty Acids," Bulletin 332, Contribution 884, Agricultural Experiment Station, University of Rhode Island, Kingston, 41 pp.) report the activity of certain fatty acids against nematodes. In 1977 Sitaramaiah and Singh (Sitaramaiah, K., R.S. Singh [1977] *Indian J. Nematol.* 7:58-65) also examined the response of nematodes to fatty acids. The results of these tests with short chain acids were equivocal, showing nematode-inhibitory action in some instances and stimulatory activity in other instances. Phytotoxicity of these acids was observed at higher concentrations. The short chain fatty acids were also examined by Malik and Jairajpuri (Malik, Z., M.S. Jairajpuri [1977] *Nematol. medit.* 12:73-79), who observed nematode toxicity at high concentrations of the fatty acids.

[0006] Notwithstanding the foregoing (some of the limitations of and problems associated with these approaches are discussed above), there is a need for safe and effective alternatives for controlling nematodes.

[0007] One method for disrupting normal cellular processes is by the use double-stranded interfering RNA (RNAi), or RNA-mediated interference (RNAi). When RNAi corresponding to a sense and antisense sequence of a target mRNA is introduced into a cell, the targeted mRNA is degraded and protein translation of that message is stopped. Although not yet fully understood, the mechanism of this post-transcriptional gene



silencing appears to be at least partially due to the generation of small RNA molecules, about 21 - 25 nucleotides in length, that correspond to the sense and antisense pieces of the RNAi introduced into the cell (Bass, B. L. [2000] "Double-stranded RNA as a template for gene silencing" *Cell* 101:235-238).

[0008] The specificity of this gene silencing mechanism appears to be extremely high, blocking expression only of targeted genes, while leaving other genes unaffected. A recent example of the use of RNAi; to inhibit genetic function in plants used *Agrobacterium tumefaciens*-mediated transformation of *Arabidopsis thaliana* (Chuang, C.-F. and E. M. Meyerowitz [2000] "Specific and heritable genetic interference by double-stranded RNA in *Arabidopsis thaliana*" *Proc. Natl. Acad. Sci. USA* 97:4985-4990). Chuang *et al.* describe the construction of vectors delivering variable levels of RNAi targeted to each of four genes involved in floral development. Severity of abnormal flower development varied between transgenic lines. For one of the genes, AGAMOUS (AG), a strong correlation existed between declining accumulation of mRNA and increasingly severe phenotypes, suggesting that AG-specific endogenous mRNA is the target of RNAi.

#### Brief Summary of the Invention

[0009] The subject invention provides novel methods and compositions for controlling nematodes. More specifically, the subject invention provides polynucleotide sequences that encode nematode genes, RNAi that selectively targets mRNA transcripts of these essential nematode genes, and methods of using these sequences in nematode control strategies. Such sequences for use according to the subject invention are summarized in Appendix 1. RNAi molecules disclosed herein can be used to inhibit the expression of one or more of these genes in nematodes.

### Brief Description of the Drawings

[00010] **Figure 1:** Modular Binary Construct System (MBCS): A series of six, 8-base cutter restriction enzyme sites has been placed between the left and right Ti borders of a previously created kan<sup>R</sup>/tet<sup>R</sup> binary plasmid.

[00011] **Figure 2:** An exemplary shuttle vector created for cloning of useful DNA fragments by containing the multi-cloning site (MCS) of a modified Bluescript plasmid flanked by 8-base restriction sites.

[00012] **Figure 3:** An exemplary shuttle vector with exemplary inserts.

[00013] **Figure 4:** A suggested RNAi binary vector with exemplary inserts.

[00014] **Figure 5:** Exemplary selectable markers for MBCS.

[00015] **Figure 6:** Exemplary scorable markers for MCBS.

[00016] **Figure 7:** Exemplary RNAi binary vector.

[00017] **Figure 8:** Exemplary RNAi shuttle vector.

### Brief Description of the Sequences

[00018] Brief Description of the Sequences can be found in Appendix I.

### Detailed Disclosure of the Invention

[00019] The subject invention provides novel methods and compositions for controlling nematodes. More specifically, the subject invention provides polynucleotide sequences and methods of using these sequences in nematode control strategies. A preferred method for controlling nematodes according to the subject invention provides materials and methods for controlling nematodes by using double-stranded interfering RNA (RNAi), or RNA-mediated interference (RNAi). The terms RNAi and RNAi are used interchangeably herein unless otherwise noted.

[00020] In one embodiment of the invention, RNAi molecules are provided which are useful in methods of killing nematodes and/or inhibiting their growth, development, parasitism or reproduction. RNAi molecules of the invention are also useful for the regulation of levels of specific mRNA in nematodes.

[00021] dsRNA (RNAi) typically comprises a polynucleotide sequence identical to a target gene (or fragment thereof) linked directly, or indirectly, to a polynucleotide

sequence complementary to the sequence of the target gene (or fragment thereof). The dsRNA may comprise a polynucleotide linker (stuffer) sequence of sufficient length to allow for the two polynucleotide sequences to fold over and hybridize to each other; however, a linker sequence is not necessary. The linker (stuffer) sequence is designed to separate the antisense and sense strands of RNAi significantly enough to limit the effects of steric hindrances and allow for the formation of dsRNA molecules.

[00022] RNA containing a nucleotide sequence identical to a fragment of the target gene is preferred for inhibition; however, RNA sequences with insertions, deletions, and point mutations relative to the target sequence can also be used for inhibition. Sequence identity may be optimized by sequence comparison and alignment algorithms known in the art (see Gribskov and Devereux, *Sequence Analysis Primer*, Stockton Press, 1991, and references cited therein) and calculating the percent difference between the nucleotide sequences by, for example, the Smith-Waterman algorithm as implemented in the BESTFIT software program using default parameters (e.g., University of Wisconsin Genetic Computing Group). Alternatively, the duplex region of the RNA may be defined functionally as a nucleotide sequence that is capable of hybridizing with a fragment of the target gene transcript.

[00023] As disclosed herein, 100% sequence identity between the RNA and the target gene is not required to practice the present invention. Thus the invention has the advantage of being able to tolerate sequence variations that might be expected due to genetic mutation, strain polymorphism, or evolutionary divergence.

[00024] RNA may be synthesized either *in vivo* or *in vitro*. Endogenous RNA polymerase of the cell may mediate transcription *in vivo*, or cloned RNA polymerase can be used for transcription *in vivo* or *in vitro*. For transcription from a transgene *in vivo* or an expression construct, a regulatory region (e.g., promoter, enhancer, silencer, splice donor and acceptor, polyadenylation) may be used to transcribe the RNA strand (or strands). Inhibition may be targeted by specific transcription in an organ, tissue, or cell type; stimulation of an environmental condition (e.g., infection, stress, temperature, chemical inducers); and/or engineering transcription at a developmental stage or age. The RNA strands may or may not be polyadenylated; the RNA strands may or may not be capable of being translated into a polypeptide by a cell's translational apparatus. RNA

may be chemically or enzymatically synthesized by manual or automated reactions. The RNA may be synthesized by a cellular RNA polymerase or a bacteriophage RNA polymerase (e.g., T3, T7, SP6). The use and production of an expression construct are known in the art (see, for example, WO 97/32016; U.S. Pat. Nos. 5,593,874; 5,698,425; 5,712,135; 5,789,214; and 5,804,693; and the references cited therein). If synthesized chemically or by *in vitro* enzymatic synthesis, the RNA may be purified prior to introduction into the cell. For example, RNA can be purified from a mixture by extraction with a solvent or resin, precipitation, electrophoresis, chromatography, or a combination thereof. Alternatively, the RNA may be used with no or a minimum of purification to avoid losses due to sample processing. The RNA may be dried for storage or dissolved in an aqueous solution. The solution may contain buffers or salts to promote annealing, and/or stabilization of the duplex strands.

[00025] Preferably and most conveniently, RNAi can be targeted to an entire polynucleotide sequence of a gene set forth herein. Preferred RNAi molecules of the instant invention are highly homologous or identical to the polynucleotides summarized in Appendix 1. The homology is preferably greater than 90% and is most preferably greater than 95%.

[00026] Fragments of genes can also be targeted. These fragments are typically in the approximate size range of about 20 nucleotides. Thus, targeted fragments are preferably at least about 15 nucleotides. In certain embodiments, the gene fragment targeted by the RNAi molecule is about 20-25 nucleotides in length. However, other size ranges can also be used. For example, using a *C. elegans* microinjection assay, RNAi "fragments" of about 60 nucleotides with between 95 and 100% identity (to a nematode gene) were determined to cause excellent inhibition.

[00027] Thus, RNAi molecules of the subject invention are not limited to those that are targeted to the full-length polynucleotide or gene. The nematode gene product can be inhibited with a RNAi molecule that is targeted to a portion or fragment of the exemplified polynucleotides; high homology (90-95%) or identity is also preferred, but not necessarily essential, for such applications.

[00028] The polynucleotide sequences identified in Appendix A and shown in the Sequence ID listing are from genes encoding nematode proteins having the functions

shown in Appendix 1. The genes exemplified herein are representative of particular classes of proteins which are preferred targets for disruption according to the subject invention. These classes of proteins include, for example, proteins involved in ribosome assembly; neuro transmitter receptors and ligands; electron transport proteins; metabolic pathway proteins; and protein and polynucleotide production, folding, and processing proteins.

[00029] Genetic regulatory sequences, such as promoters, enhancers, and terminators, can be used in genetic constructs to practice the subject invention. Such constructs themselves can also be used for nematode control. Various constructs can be used to achieve expression in specific plant tissues (by using root specific promoters, for example) and/or to target specific nematode tissues (by using targeting elements or adjacent targeting sequences, for example).

[00030] In a specific embodiment of the subject invention, plant cells, preferably root cells, are genetically modified to produce at least one RNAi that is designed to be taken up by nematodes during feeding to block expression (or the function of) of a target gene. As is known in the art, RNAi can target and reduce (and, in some cases, prevent) the translation of a specific gene product. RNAi can be used to reduce or prevent message translation in any tissue of the nematode because of its ability to cross tissue and cellular boundaries. Thus, RNAi that is contacted with a nematode by soaking, injection, or consumption of a food source will cross tissue and cellular boundaries. RNAi can also be used as an epigenetic factor to prevent the proliferation of subsequent generations of nematodes.

[00031] Nematode polynucleotide sequences disclosed herein demonstrate conserved nucleotide motifs among different nematode genera. Conserved nucleotide motifs strongly suggest that these sequences are associated with viability and/or parasitism and are functionally conserved and expressed in both *Meloidogyne incognita* (root-knot nematode) and *Globodera rostochiensis* and *Globodera pallida* (potato cyst nematodes). The use of these polynucleotides, and RNAi inhibitors thereof, is advantageous because such RNAi can be designed to have broad RNAi specificity and are thus useful for controlling a large number of plant parasitic nematodes *in planta*. Because the genes identified in this disclosure are associated with nematode survival

and/or parasitism, RNAi inhibition of these genes (arising from contacting nematodes with compositions comprising RNAi molecules) prevents and/or reduces parasitic nematode growth, development, and or parasitism.

[00032] Methods of the subject invention include the transformation of plant cells with genes or polynucleotides of the present invention, which can be used to produce nematode inhibitors or RNAi in the plants. In one embodiment, the transformed plant or plant tissue can express RNAi molecules encoded by the gene or polynucleotide sequence introduced into the plant. Other nematode inhibitors contemplated by the invention include antisense molecules specific to the polynucleotide sequences disclosed herein. The transformation of plants with genetic constructs disclosed herein can be accomplished using techniques well known to those skilled in the art and can involve modification of the gene(s) to optimize expression in the plant to be made resistant to nematode infection and infestation. Furthermore, it is known in the art that many tissues of the transgenic plants (such as the roots) can be targeted for transformation.

[00033] RNA-mediated interference (RNAi) of gene expression. Several aspects of root-knot nematode biology make classical genetic studies difficult with this organism. Since root-knot nematodes reproduce by obligatory mitotic parthenogenesis, the opportunity to perform genetic crosses is not available. Microinjection of RNAi can be used to manipulate gene expression in *C. elegans* (Fire, A., S. Xu, M. K. Montgomery, S. A. Kostas, S. E. Driver, and C. C. Mello. [1998] "Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*" *Nature* 391:806-811). Microinjecting (into adult nematodes) RNAi can turn off specific genes in progeny worms complementary to the coding region of the genes. Moreover, gene inhibition occurs in progeny when RNAi is injected into the body cavity of the adult, indicating the ability of the RNAi to cross cellular boundaries. This RNAi injection method provides a molecular genetic tool that allows for analysis of gene function in root-knot nematodes.

[00034] RNAi can be taken up by *C. elegans* by simply soaking the nematodes in a solution RNAi. This results in targeted inhibition of gene expression in the nematode (Maeda, I., Y. Kohara, M. Yamamoto and A. Sugimoto [1999] "RNAi screening with a non-redundant cDNA set" International Worm Meeting, Madison, WI, abstract 565). Nematodes fed *E. coli* expressing RNAi also demonstrate targeted and

heritable inhibition of gene expression (Sarkissian, M., H. Tabara and C. C. Mello [1999] "A mut-6 screen for RNAi deficient mutants" International Worm Meeting, Madison, WI, abstract 741; Timmons, L. and A. Fire [1998] "Specific interference by ingested dsRNA" *Nature* 395:854; WO 99/32619, hereby incorporated by reference in its entirety).

[00035] Accordingly, one aspect of the instant invention is directed to the control of nematodes comprising contacting nematodes with compositions comprising RNAi molecules specific to the nematode genes disclosed herein. The contacting step may include soaking the nematodes in a solution containing RNAi molecules, feeding nematodes RNAi molecules contained in microbes or plant cells upon which the nematode feeds, or injecting nematodes with RNAi. Nematodes can also be "contacted" and controlled by RNAi expressed in plant tissues that would be consumed, ingested, or frequented by nematodes.

[00036] The RNAi molecules provided to the nematodes may be specific to a single gene. A "cocktail" of RNAi molecules specific to various segments of a single gene can also be used. In addition, a "multigene cocktail" of RNAi molecules specific to two or more genes (or segments thereof) may be applied to the nematodes according to the subject invention.

[00037] In addition to RNAi uptake mediated by transgenic plants, nematodes can be directly transformed with RNAi constructs of cDNAs encoding secretory or other essential proteins to reduce expression of the corresponding gene. The transgenic animals can be assayed for inhibition of gene product using immunoassays or for reduced virulence on a host. Progeny of affected worms can also be assayed by similar methods.

[00038] Procedures that can be used for the preparation and injection of RNAi include those detailed by Fire *et al.*, (1998; <ftp://ciw1.ciwemb.edu>). Root-knot nematodes can be routinely monoxenically cultured on *Arabidopsis thaliana* roots growing on Gamborg's B-5/Gelrite® media. This nematode-host pathosystem is ideally suited for these microinjection experiments since limited root galling results in the parasitic stages (late J2 through adult females) developing outside of the root for easy accessibility for injecting. Another advantage is the parthenogenic reproduction of root-knot nematodes, which makes fertilization by males unnecessary for egg production. The RNAi can be injected into the body cavity of parasitic stages of root-knot nematodes

feeding on *A. thaliana* roots using microinjection. Control nematodes can be injected in parallel with only buffer or an unrelated RNAi. Injected nematodes can be monitored for egg production, and the eggs can be collected for the assays described below. Female root-knot nematodes will typically survive and lay more than 250 eggs following 1  $\mu$ l injection of buffer.

[00039] Alternatively, methods are available for microinjecting materials directly into the plant root cells upon which nematodes feed: giant cells or syncytial cells (Böckenhoff, A. and F.M.W. Grundler [1994] "Studies on the nutrient uptake by the beet cyst nematode *Heterodera schachtii* by *in situ* microinjection of fluorescent probes into the feeding structures in *Arabidopsis thaliana*" *Parasitology* 109:249-254). This provides an excellent test system to screen RNAi molecules for efficacy by directly inhibiting growth and development of the nematode feeding upon the microinjected plant cell, or by reducing fecundity and the ability of said nematode to generate pathogenic or viable progeny.

[00040] There are a number of strategies that can be followed to assay for RNAi gene interference. Inhibition of gene expression by RNAi inhibits the accumulation of the corresponding secretory protein in the esophageal gland cells of transgenic J2 hatched from the eggs produced by the injected nematodes. In the first assay, polyclonal antibodies to the target gene product can be used in immunolocalization studies (Hussey, R. S. [1989] "Monoclonal antibodies to secretory granules in esophageal glands of *Meloidogyne* species" *J. Nematol.* 21:392-398; Borgonie, G, E. van Driessche, C. D. Link, D. de Waele, and A. Coomans [1994] "Tissue treatment for whole mount internal lectin staining in the nematodes *Caenorhabditis elegans*, *Panagrolaimus superbus* and *Acrobeloides maximus*" *Histochemistry* 101:379-384) to monitor the synthesis of the target protein in the gland cells of progeny of the injected nematodes, or in any other nematode tissue that fails to express the essential targeted gene. Interference of endogenous gene activity by the RNAi eliminates binding of the antibodies to secretory granules in the glands, or any other target tissue, of the transgenic nematodes, and can be monitored by these *in situ* hybridization experiments. Control nematodes injected only with the injection buffer can be processed similar to the RNAi treated nematodes.



[00041] Another assay is designed to determine the effect of the RNAi on reducing the virulence of J2 progeny of the injected females. Egg masses from injected females can be transferred singly to *A. thaliana* plates to assess the ability of the transgenic J2 to infect roots. The J2 hatching from the eggs transferred to the plates can be monitored; after 25 days the number of galls with egg laying females can be recorded. The *A. thaliana* roots can also be stained with acid fuchsin to enumerate the number of nematodes in the roots. Egg masses from nematodes injected only with the injection buffer can be handled similarly and used as controls. The treatments can be replicated, and the root infection data can be analyzed statistically. These experiments can be used to assess the importance of the target genes in root-knot nematode's virulence or viability. By staining the J2 progeny of the injected females with the antibodies, it can be determined whether RNAi blocks expression of the targeted gene.

[00042] Additional uses of polynucleotides. The polynucleotide sequences exemplified herein can be used in a variety of ways. These polynucleotides can be used in assays for additional polynucleotides and additional homologous genes, and can be used in tracking the quantitative and temporal expression of parasitism genes in nematodes. These polynucleotides can be cloned into microbes for production and isolation of their gene products. Among the many uses of the isolated gene product is the development of additional inhibitors and modifiers. The protein products of the subject polynucleotides can also be used as diagnostic tools. For example, proteins encoded by the parasitism genes, as identified herein, can be used in large scale screenings for additional peptide inhibitors. The use of peptide phage display screening is one method that can be used in this regard. Thus, the subject invention also provides new biotechnological strategies for managing nematodes under sustainable agricultural conditions.

[00043] Antisense technologies can also be used for phytopathogenic nematode control. Antisense technology can be used to interfere with expression of the disclosed endogenous nematode genes. Antisense technology can also be used to alter the components of plants used as targets by the nematodes. For example, the transformation of a plant with the reverse complement of an endogenous gene encoded by a polynucleotide exemplified herein can result in strand co-suppression and gene silencing

or inhibition of a target involved in the nematode infection process. Thus, the subject invention includes transgenic plants (which are preferably made nematode-resistant in this manner, and other organisms including microbes and phages) comprising RNAi or antisense molecules specific to any of the polynucleotides identified herein.

[00044] Polynucleotide probes. DNA possesses a fundamental property called base complementarity. In nature, DNA ordinarily exists in the form of pairs of anti-parallel strands, the bases on each strand projecting from that strand toward the opposite strand. The base adenine (A) on one strand will always be opposed to the base thymine (T) on the other strand, and the base guanine (G) will be opposed to the base cytosine (C). The bases are held in apposition by their ability to hydrogen bond in this specific way. Though each individual bond is relatively weak, the net effect of many adjacent hydrogen bonded bases, together with base stacking effects, is a stable joining of the two complementary strands. These bonds can be broken by treatments such as high pH or high temperature, and these conditions result in the dissociation, or "denaturation," of the two strands. If the DNA is then placed in conditions which make hydrogen bonding of the bases thermodynamically favorable, the DNA strands will anneal, or "hybridize," and reform the original double-stranded DNA. If carried out under appropriate conditions, this hybridization can be highly specific. That is, only strands with a high degree of base complementarity will be able to form stable double-stranded structures. The relationship of the specificity of hybridization to reaction conditions is well known. Thus, hybridization may be used to test whether two pieces of DNA are complementary in their base sequences. It is this hybridization mechanism which facilitates the use of probes of the subject invention to readily detect and characterize DNA sequences of interest.

[00045] The specifically exemplified polynucleotides of the subject invention can themselves be used as probes. Additional polynucleotide sequences can be added to the ends of (or internally in) the exemplified polynucleotide sequences so that polynucleotides that are longer than the exemplified polynucleotides can also be used as probes. Thus, isolated polynucleotides comprising one or more of the exemplified sequences are within the scope of the subject invention. Polynucleotides that have less nucleotides than the exemplified polynucleotides can also be used and are contemplated within the scope of the present invention. For example, for some purposes, it might be

useful to use a conserved sequence from an exemplified polynucleotide wherein the conserved sequence comprises a portion of an exemplified sequence. Thus, polynucleotides of the subject invention can be used to find additional, homologous (wholly or partially) genes.

[00046] Probes of the subject invention may be composed of DNA, RNA, or PNA (peptide nucleic acid). The probe will normally have at least about 10 bases, more usually at least about 17 bases, and may have about 100 bases or more. Longer probes can readily be utilized, and such probes can be, for example, several kilobases in length. The probe sequence is designed to be at least substantially complementary to a portion of a gene encoding a protein of interest. The probe need not have perfect complementarity to the sequence to which it hybridizes. The probes may be labeled utilizing techniques that are well known to those skilled in this art.

[00047] One approach for the use of the subject invention as probes entails first identifying DNA segments that are homologous with the disclosed nucleotide sequences using, for example, Southern blot analysis of a gene bank. Thus, it is possible, without the aid of biological analysis, to know in advance the probable activity of many new polynucleotides, and of the individual gene products expressed by a given polynucleotide. Such an analysis provides a rapid method for identifying commercially valuable compositions.

[00048] One hybridization procedure useful according to the subject invention typically includes the initial steps of isolating the DNA sample of interest and purifying it chemically. Either lysed nematodes or total fractionated nucleic acid isolated from nematodes can be used. Cells can be treated using known techniques to liberate their DNA (and/or RNA). The DNA sample can be cut into pieces with an appropriate restriction enzyme. The pieces can be separated by size through electrophoresis in a gel, usually agarose or acrylamide. The pieces of interest can be transferred to an immobilizing membrane.

[00049] The particular hybridization technique is not essential to the subject invention. As improvements are made in hybridization techniques, they can be readily applied.

[00050] The probe and sample can then be combined in a hybridization buffer solution and held at an appropriate temperature until annealing occurs. Thereafter, the membrane is washed free of extraneous materials, leaving the sample and bound probe molecules typically detected and quantified by autoradiography and/or liquid scintillation counting. As is well known in the art, if the probe molecule and nucleic acid sample hybridize by forming a strong non-covalent bond between the two molecules, it can be reasonably assumed that the probe and sample are essentially identical or very similar. The probe's detectable label provides a means for determining in a known manner whether hybridization has occurred.

[00051] In the use of the nucleotide segments as probes, the particular probe is labeled with any suitable label known to those skilled in the art, including radioactive and non-radioactive labels. Typical radioactive labels include  $^{32}\text{P}$ ,  $^{35}\text{S}$ , or the like. Non-radioactive labels include, for example, ligands such as biotin or thyroxine, as well as enzymes such as hydrolases or peroxidases, or the various chemiluminescers such as luciferin, or fluorescent compounds like fluorescein and its derivatives. In addition, the probes can be made inherently fluorescent as described in International Application No. WO 93/16094.

[00052] Various degrees of stringency of hybridization can be employed. The more stringent the conditions, the greater the complementarity that is required for duplex formation. Stringency can be controlled by temperature, probe concentration, probe length, ionic strength, time, and the like. Preferably, hybridization is conducted under moderate to high stringency conditions by techniques well known in the art, as described, for example, in Keller, G.H., M.M. Manak (1987) *DNA Probes*, Stockton Press, New York, NY., pp. 169-170.

[00053] As used herein "moderate to high stringency" conditions for hybridization refers to conditions that achieve the same, or about the same, degree of specificity of hybridization as the conditions "as described herein." Examples of moderate to high stringency conditions are provided herein. Specifically, hybridization of immobilized DNA on Southern blots with  $^{32}\text{P}$ -labeled gene-specific probes was performed using standard methods (Maniatis *et al.*). In general, hybridization and subsequent washes were carried out under moderate to high stringency conditions that

allowed for detection of target sequences with homology to sequences exemplified herein. For double-stranded DNA gene probes, hybridization was carried out overnight at 20-25° C below the melting temperature ( $T_m$ ) of the DNA hybrid in 6X SSPE, 5X Denhardt's solution, 0.1% SDS, 0.1 mg/ml denatured DNA. The melting temperature is described by the following formula from Beltz *et al.* (1983):

[00054]  $T_m = 81.5^\circ\text{C} + 16.6 \log[\text{Na}^+] + 0.41(\%G+C) - 0.61(\%\text{formamide}) - 600/\text{length of duplex in base pairs}.$

Washes are typically carried out as follows:

- (1) Twice at room temperature for 15 minutes in 1X SSPE, 0.1% SDS (low stringency wash).
- (2) Once at  $T_m - 20^\circ\text{C}$  for 15 minutes in 0.2X SSPE, 0.1% SDS (moderate stringency wash).

[00055] For oligonucleotide probes, hybridization was carried out overnight at 10-20°C below the melting temperature ( $T_m$ ) of the hybrid in 6X SSPE, 5X Denhardt's solution, 0.1% SDS, 0.1 mg/ml denatured DNA.  $T_m$  for oligonucleotide probes was determined by the following formula from Suggs *et al.* (1981):

[00056]  $T_m (^\circ\text{C}) = 2(\text{number T/A base pairs}) + 4(\text{number G/C base pairs})$

[00057] Washes were typically carried out as follows:

- [00058] (1) Twice at room temperature for 15 minutes 1X SSPE, 0.1% SDS (low stringency wash).
- [00059] (2) Once at the hybridization temperature for 15 minutes in 1X SSPE, 0.1% SDS (moderate stringency wash).

[00060] In general, salt and/or temperature can be altered to change stringency. With a labeled DNA fragment of greater than about 70 or so bases in length, the following conditions can be used:

Low:	1 or 2X SSPE, room temperature
Low:	1 or 2X SSPE, 42°C
Moderate:	0.2X or 1X SSPE, 65°C
High:	0.1X SSPE, 65°C.

[00061] Duplex formation and stability depend on substantial complementarity between the two strands of a hybrid, and, as noted above, a certain degree of mismatch

can be tolerated. Therefore, polynucleotide sequences of the subject invention include mutations (both single and multiple), deletions, and insertions in the described sequences, and combinations thereof, wherein said mutations, insertions, and deletions permit formation of stable hybrids with a target polynucleotide of interest. Mutations, insertions, and deletions can be produced in a given polynucleotide sequence using standard methods known in the art. Other methods may become known in the future.

[00062] The mutational, insertional, and deletional variants of the polynucleotide sequences of the invention can be used in the same manner as the exemplified polynucleotide sequences so long as the variants have substantial sequence similarity with the original sequence. As used herein, substantial sequence similarity refers to the extent of nucleotide similarity that is sufficient to enable the variant polynucleotide to function in the same capacity as the original sequence. Preferably, this similarity is greater than 50%; more preferably, this similarity is greater than 75%; and most preferably, this similarity is greater than 90%. The degree of similarity needed for the variant to function in its intended capacity will depend upon the intended use of the sequence. It is well within the skill of a person trained in this art to make mutational, insertional, and deletional mutations that are designed to improve the function of the sequence or otherwise provide a methodological advantage.

[00063] PCR technology. Polymerase Chain Reaction (PCR) is a repetitive, enzymatic, primed synthesis of a nucleic acid sequence. This procedure is well known and commonly used by those skilled in this art (see U.S. Patent Nos. 4,683,195; 4,683,202; and 4,800,159; Saiki *et al.*, 1985). PCR is based on the enzymatic amplification of a DNA fragment of interest that is flanked by two oligonucleotide primers that hybridize to opposite strands of the target sequence. The primers are oriented with the 3' ends pointing towards each other. Repeated cycles of heat denaturation of the template, annealing of the primers to their complementary sequences, and extension of the annealed primers with a DNA polymerase result in the amplification of the segment defined by the 5' ends of the PCR primers. Since the extension product of each primer can serve as a template for the other primer, each cycle essentially doubles the amount of DNA fragment produced in the previous cycle. This results in the exponential accumulation of the specific target fragment, up to several million-fold in a

few hours. By using a thermostable DNA polymerase such as *Taq* polymerase, which is isolated from the thermophilic bacterium *Thermus aquaticus*, the amplification process can be completely automated. Other enzymes that can be used are known to those skilled in the art.

[00064] The polynucleotide sequences of the subject invention (and portions thereof such as conserved regions and portions that serve to distinguish these sequences from previously-known sequences) can be used as, and/or used in the design of, primers for PCR amplification. In performing PCR amplification, a certain degree of mismatch can be tolerated between primer and template. Therefore, mutations, deletions, and insertions (especially additions of nucleotides to the 5' end) of the exemplified polynucleotides can be used in this manner. Mutations, insertions and deletions can be produced in a given primer by methods known to an ordinarily skilled artisan.

[00065] The polynucleotide sequences of the instant invention may be "operably linked" to regulatory sequences such as promoters and enhancers. Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is "operably linked" to DNA encoding a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is "operably linked" to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is "operably linked" to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

[00066] Polynucleotides and proteins. Polynucleotides of the subject invention can be defined according to several parameters. One characteristic is the biological activity of the protein products as identified herein. The proteins and genes of the subject invention can be further defined by their amino acid and nucleotide sequences. The sequences of the molecules can be defined in terms of homology to certain exemplified sequences as well as in terms of the ability to hybridize with, or be amplified by, certain

exemplified probes and primers. Additional primers and probes can readily be constructed by those skilled in the art such that alternate polynucleotide sequences encoding the same amino acid sequences can be used to identify and/or characterize additional genes. The proteins of the subject invention can also be identified based on their immunoreactivity with certain antibodies.

[00067] The polynucleotides and proteins of the subject invention include portions, fragments, variants, and mutants of the full-length sequences as well as fusions and chimerics, so long as the encoded protein retains the characteristic biological activity of the proteins identified herein. As used herein, the terms "variants" or "variations" of genes refer to nucleotide sequences that encode the same proteins or which encode equivalent proteins having equivalent biological activity. As used herein, the term "equivalent proteins" refers to proteins having the same or essentially the same biological activity as the exemplified proteins.

[00068] It will be apparent to a person skilled in this art that genes within the scope of the subject invention can be identified and obtained through several means. The specific genes exemplified herein may be obtained from root-knot nematodes. Genes, or portions or variants thereof, may also be artificially synthesized by, for example, a gene synthesizer.

[00069] Variations of genes may be readily constructed using standard techniques such as site-directed mutagenesis and other methods of making point mutations and by DNA shuffling, for example. In addition, gene and protein fragments can be made using commercially available exonucleases, endonucleases, and proteases according to standard procedures. For example, enzymes such as *Bal31* can be used to systematically cut off nucleotides from the ends of genes. In addition, genes that encode fragments may be obtained using a variety of restriction enzymes. Proteases may be used to directly obtain active fragments of these proteins. Of course, molecular techniques for cloning polynucleotides and producing gene constructs of interest are also well known in the art. *In vitro* evaluation techniques, such as MAXYGEN's "Molecular Breeding" can also be applied to practice the subject invention.

[00070] Other molecular techniques can also be applied using the teachings provided herein. For example, antibodies raised against proteins encoded by



polynucleotides disclosed herein can be used to identify and isolate proteins from a mixture of proteins. Specifically, antibodies may be raised to the portions of the proteins that are conserved and most distinct from other proteins. These antibodies can then be used to specifically identify equivalent proteins by immunoprecipitation, enzyme linked immunosorbent assay (ELISA), or Western blotting. Antibodies to proteins encoded by polynucleotides disclosed herein, or to equivalent proteins, can readily be prepared using standard procedures known in the art. The genes that encode these proteins can be obtained from various organisms.

[00071] Because of the redundancy of the genetic code, a variety of different DNA sequences can encode the amino acid sequences encoded by the polynucleotide sequences disclosed herein. It is well within the skill of a person trained in the art to create these alternative DNA sequences encoding proteins having the same, or essentially the same, amino acid sequence. These variant DNA sequences are within the scope of the subject invention. As used herein, reference to "essentially the same" sequence refers to sequences that have amino acid substitutions, deletions, additions, or insertions that do not materially affect biological activity. Fragments retaining the characteristic biological activity are also included in this definition.

[00072] A further method for identifying genes and polynucleotides (and the proteins encoded thereby) of the subject invention is through the use of oligonucleotide probes. Probes provide a rapid method for identifying genes of the subject invention. The nucleotide segments that are used as probes according to the invention can be synthesized using a DNA synthesizer and standard procedures.

[00073] The subject invention comprises variant or equivalent proteins (and nucleotide sequences coding for equivalent proteins or for inhibitors of the genes encoding such proteins) having the same or similar biological activity of inhibitors or proteins encoded by the exemplified polynucleotides. Equivalent proteins will have amino acid similarity with an exemplified protein (or peptide). The amino acid and/or nucleotide identity will typically be greater than 60%. Preferably, the identity will be greater than 75%. More preferably, the identity will be greater than 80%, and even more preferably greater than 90%. Most preferably, the identity will be greater than 95%. RNAi molecules will also have corresponding identities in these preferred ranges. These

identities are as determined using standard alignment techniques for determining amino acid and/or nucleotide identity. The identity/similarity will be highest in critical regions of the protein or gene including those regions that account for biological activity or that are involved in the determination of three-dimensional configuration that is ultimately responsible for the biological activity. In this regard, certain amino acid substitutions are acceptable and can be expected if these substitutions are in regions which are not critical to activity or are conservative amino acid substitutions which do not affect the three-dimensional configuration of the molecule. For example, amino acids may be placed in the following classes: non-polar, uncharged polar, basic, and acidic. Conservative substitutions whereby an amino acid of one class is replaced with another amino acid of the same type fall within the scope of the subject invention so long as the substitution does not materially alter the biological activity of the compound. Below is a list of examples of amino acids belonging to various classes

Class of Amino Acid	Examples of Amino Acids
Nonpolar	Ala, Val, Leu, Ile, Pro, Met, Phe, Trp
Uncharged Polar	Gly, Ser, Thr, Cys, Tyr, Asn, Gln
Acidic	Asp, Glu
Basic	Lys, Arg, His

[00074] In some instances, non-conservative substitutions can also be made. The critical factor is that these substitutions must not detract from the ability to manage nematode-caused diseases.

[00075] An "isolated" or "substantially pure" nucleic acid molecule or polynucleotide is a polynucleotide that is substantially separated from other polynucleotide sequences which naturally accompany a nucleic acid molecule. The term embraces a polynucleotide sequence which was removed from its naturally occurring environment by the hand of man. This includes recombinant or cloned DNA isolates,

chemically synthesized analogues and analogues biologically synthesized by heterologous systems. An "isolated" or "purified" protein, likewise, is a protein removed from its naturally occurring environment.

[00076] Recombinant hosts. The genes, antisense, and RNAi polynucleotides within the scope of the present invention can be introduced into a wide variety of microbial or plant hosts. Plant cells can be transformed (made recombinant) in this manner. Microbes, for example, can also be used in the application of RNAi molecules of the subject invention in view of the fact that microbes are a food source for nematodes

[00077] There are many methods for introducing a heterologous gene or polynucleotide into a host cell or cells under conditions that allow for stable maintenance and expression of the gene or polynucleotide. These methods are well known to those skilled in the art. Synthetic genes, such as, for example, those genes modified to enhance expression in a heterologous host (such as by preferred codon usage or by the use of adjoining, downstream, or upstream enhancers) that are functionally equivalent to the genes (and which encode equivalent proteins) can also be used to transform hosts. Methods for the production of synthetic genes are known in the art.

[00078] Where the gene or polynucleotide of interest is introduced via a suitable vector into a microbial host, and said host is applied to the environment in a living state, certain host microbes are preferred. Certain microorganism hosts are known to occupy the phytosphere, phylloplane, phyllosphere, rhizosphere, and/or rhizoplane of one or more crops of interest. These microorganisms can be selected so as to be capable of successfully competing in the particular environment (crop and other habitats) with the wild-type microorganisms, provide for stable maintenance and expression of the gene expressing a polypeptide of interest, and, desirably, provide for improved protection of the protein/peptide from environmental degradation and inactivation.

[00079] A large number of microorganisms is known to inhabit the phylloplane (the surface of the plant leaves) and/or the rhizosphere (the soil surrounding plant roots) of a wide variety of important crops. These microorganisms include bacteria, algae, and fungi. Of particular interest are microorganisms, such as bacteria, *e.g.*, genera *Pseudomonas*, *Erwinia*, *Serratia*, *Klebsiella*, *Xanthomonas*, *Streptomyces*, *Rhizobium*, *Rhodopseudomonas*, *Methylophilus*, *Agrobacterium*, *Acetobacter*, *Lactobacillus*,

*Arthrobacter*, *Azotobacter*, *Leuconostoc*, and *Alcaligenes*; fungi, particularly yeast, *e.g.*, genera *Saccharomyces*, *Cryptococcus*, *Kluyveromyces*, *Sporobolomyces*, *Rhodotorula*, and *Aureobasidium*. Of particular interest are the pigmented microorganisms.

[00080] Methods of the subject invention also include the transformation of plants or plant tissue with genes which encode the RNAi molecules of the present invention. In one embodiment, the transformed plant or plant tissue expresses antisense RNA and/or RNAi. Transformation of cells can be made by those skilled in the art using standard techniques. Materials necessary for these transformations are disclosed herein or are otherwise readily available to the skilled artisan.

[00081] Additional methods and formulations for control of pests. Control of nematode pests using the RNAi molecules of the instant invention can be accomplished by a variety of additional methods that would be apparent to those skilled in the art having the benefit of the subject disclosure. A "cocktail" of two or more RNAi molecules can be used to disrupt one or more of the genes identified herein. The "cocktail" of RNAi molecules may be specific to segments of a single gene or the entire gene. A "multigene cocktail" of RNAi molecules specific to two or more genes (or segments thereof) is also encompassed by the instant invention. In another embodiment of the instant invention, the disclosed RNAi molecules, cocktails, and/or multigene cocktails thereof, may be used in conjunction with other known nematode control agents and methodologies. Such cocktails can be used to combat the development of resistance by nematodes to a certain inhibitor or inhibitors.

[00082] Compositions of the subject invention which comprise RNAi molecules and carriers can be applied, themselves, directly or indirectly, to locations frequented by, or expected to be frequented by, nematodes. Microbial hosts which were transformed with polynucleotides that encode RNAi molecules, express said RNAi molecules, and which colonize roots (*e.g.*, *Pseudomonas*, *Bacillus*, and other genera) can be applied to the sites of the pest, where they will proliferate and be ingested. The result is control of the pest. Thus, methods of the subject invention include, for example, the application of recombinant microbes to the pests (or their locations). The recombinant microbes may also be transformed with more than one RNAi molecule thereby delivering a "cocktail" of RNAi molecules to the nematode pests. A carrier may be any substance suitable for

- [00089] 1. Grow *Agrobacterium* overnight in 5 mls LB + antibiotics at 30°C on shaker (for *Agrobacterium rhizogenes* strain K599 no antibiotics are needed).
- [00090] 2. Use the 5 mls of overnight culture to inoculate 500 mls LB + antibiotics at 30°C on shaker. Grow overnight.
- [00091] 3. Add liquid culture in eight 50 ml polypropylene orange cap tubes.
- [00092] 4. Centrifuge 10 min., 4000 rpm, 4°C.
- [00093] 5. Resuspend cells in each tube with 20 mls 10% glycerol (on ice)
- [00094] 6. Centrifuge 10 min., 4000 rpm, 4°C.
- [00095] 7. Resuspend cells in each tube with 10 mls 10% glycerol (on ice).
- [00096] 8. Centrifuge 10 min., 4000 rpm, 4°C.
- [00097] 9. Resuspend cells in each tube with 2 mls 10% glycerol (on ice).
- [00098] 10. Aliquot 50 µl into cold Eppendorf tube and place onto dry ice.
- [00099] 11. Store electro-competent cells at -80°C. These cells can be used for up to two years.

[000100] Electroporations:

- [000101] 1. Add 1 µl to 5 µl of DNA (resuspended in H<sub>2</sub>O and not TE or other buffer) to 50 µl of *Agrobacterium* electrocompetent cells and mix.
- [000102] 2. Transfer 20 µl of DNA/*Agrobacterium* mix to cuvette.
- [000103] 3. Electroporate:  
25µF, 400 Ω resistance, 2.5 volts (0.2cm cuvette) or 1.8 volts (0.1cm cuvette for BioRad electroporator. 330 µF, 4000 kΩ, low w, fast charge rate for BRL Electroporator.
- [000104] 4. Add 1ml of LB and transfer to Eppendorf tube.
- [000105] 5. Shake at 30°C for 2 hours.
- [000106] 6. Centrifuge down cells (2 min. 14 krpm).
- [000107] 7. Plate all onto LB + antibiotics (most *Agrobacterium* strains are naturally streptomycin resistant).

Example 3 – Protocol for Production of Transgenic Hairy Roots on Soybean

delivering the RNAi molecules to the nematode. Acceptable carriers are well known in the art and also are commercially available. For example, such acceptable carriers are described in E.W. Martin's *Remington's Pharmaceutical Science*, Mack Publishing Company, Easton, PA.

[00083] All patents, patent applications, provisional applications, and publications referred to or cited herein are incorporated by reference in their entirety to the extent they are not inconsistent with the explicit teachings of this specification.

[00084] Following are examples that illustrate procedures for practicing the invention. These examples should not be construed as limiting. All percentages are by weight and all solvent mixture proportions are by volume unless otherwise noted.

#### Example 1— Production of Hairy Roots for RNAi Testing

[00085] A hairy root assay system was developed for testing the anti-nematode activity of RNAi molecules.

[00086] *Agrobacterium rhizogenes*: Several *Agrobacterium rhizogenes* strains produce hairy roots on a variety of plant species. *A. rhizogenes* strains, A4, 15834, 8196 and LBA4404 demonstrate hairy root development on tomato and sugar beet, with A4 being the most efficient. The *A. rhizogenes* strain K599 demonstrated very efficient formation on transgenic soybean hairy roots and was also effective on sugar beet and *Arabidopsis*. However, strain K599 failed to produce hairy roots on tomato tissues possibly due to hyper-virulence.

[00087] Hairy root production: Transgenic hairy roots were identified by stable GUS expression in tomato, sugar beet, soybean and *Arabidopsis*. The construct pAKK1401 (pNOS / NPT-II / tNOS // pSU / GUS / tNOS) was used to produce hairy roots when transformed into *A. rhizogenes* strains A4 or K599. Transgenic roots were identified by GUS expression.

#### Example 2 – Protocol for Electro-competent *Agrobacterium* and Electroporation

[00088] Electro-competent *Agrobacterium* Protocol:

[000108] Seed Sterilization. Rinse the soybean seed with 70% ETOH for 2-5 min. Remove and add 20% Clorox and shake for 20-25 min. Rinse 3X with sterile water. Plate the seed, 5 seed per plate, onto ½ MSB5 + 2% sucrose + 0.2% gel (referred to as ½ MSB5). Place seed into chamber at 25°C, 16/8 photoperiod for 5-7 day (depending on genotype) germination period. After 1 week seedlings can be placed into cold room for longer storage if necessary (not to exceed 2 weeks).

[000109] Agrobacterium Preparation. For *Agrobacterium rhizogenes* strain K599, take a small sample from frozen glycerol into 25-50 ml of NZYM media with 50 mg/L kanamycin in a 125-250 ml Erlenmeyer flask. Place onto shaker at 28-30 °C for 16 - 20 hours. Pour sample into centrifuge tube and centrifuge the bacterium at 4000 rpm for 10 min. Pour off supernatant and re-suspend the pellet with an equal volume of liquid ½ MSB5 + 200 µM acetosyringone. Use pipette to re-suspend the pellet and homogenize the sample (remove all clumps). To determine O.D., prepare a 1:10 dilution by putting 900 µl ½ MSB5 into cuvette and add 100 µl of bacterial sample. Determine the O.D.<sub>660</sub> and calculate the volume needed to adjust (dilute) OD to approximately 0.2 for inoculation. Check final O.D.

[000110] Explant Preparation and inoculation. Place a sterile filter paper onto plates of 1/2 MSB5. Cut soybean cotyledons just above the shoot apex and place onto plate. Lightly scar the cotyledon's abaxial surface (flat side, upper surface that reaches toward sun) with a scalpel blade. Cut each cotyledon transversely into 2-3 pieces (no smaller than 1 cm). Add approximately 10 ml of prepared bacterial solution to each plate and allow cotyledons to incubate for 1 hr. Remove the bacteria using a vacuum aspirator fitted with sterile pipette tip, ensure that there is no standing liquid. Orient all explants with abaxial surface up and wrap plates for a 3 day co-culture, 25°C in light (16/8 photoperiod).

[000111] Hairy root selection and maintenance. After 3 day co-culture, wash explants with liquid ½ MSB5 + 500 mg/L carbenicillin. Transfer the explants abaxial side up to selection media, ½ MSB5 supplemented with 500 mg/L carbenicillin and 200 mg/L kanamycin. Roots should develop in approximately 2-3 weeks. The roots will form primarily from the cut vascular bundles with other roots developing from the small cuts on cotyledon surface. Remove roots (>1 cm in length) and place onto replica media with

transfers to fresh media every 2 weeks to prevent *Agrobacterium* overgrowth. After 6-8 weeks on selection the roots can be moved to media without kanamycin, however carbenicillin must remain in media for several months for continued suppression of *Agrobacterium*. At this stage roots can be used for testing RNAi for nematode control. Sterilized nematodes can be added and observed for RNAi affects.

Example 4 – Testing of RNAi for Plant Parasitic Nematode Control.

[000112] Various types of nematodes can be used in appropriate bioassays. For example, *Caenorhabditis elegans*, a bacterial feeding nematode, and plant parasitic nematodes can be used for bioassay purposes. Examples of plant parasitic nematodes include a migratory endo-parasite, *Pratylenchus scribneri* (lesion), and two sedentary endo-parasites, *Meloidogyne javanica* (root-knot) and *Heterodera schachtii* (cyst).

[000113] *C. elegans*: RNAi vectors can be tested through expression of the RNAi in *E. coli*. *C. elegans* are fed *E. coli* and assayed for their growth by measuring growth of nematodes, production of eggs and viability of offspring. Another approach is to inject dsRNA directly into living nematodes. Finally, soaking nematodes in a solution of *in vitro*-prepared RNAi can quickly establish efficacy of treatment.

[000114] *P. scribneri*: The *P. scribneri in vitro* feeding assay uses a corn root exudate (CRE) as a feeding stimulus and both the red dye Amaranth or potassium, arsenate as feeding indicators. Feeding is confirmed after seven days by the presence of red stained intestinal cells in live worms exposed to the Amaranth or death of worms exposed to arsenate. This bioassay is used to test soluble toxins or RNAi. *P. scribneri* has also been cultured on wild type roots of corn, rice and *Arabidopsis*, and on A. rhizogenes-induced hairy roots of sugar beet and tomato. *P. scribneri* is very valuable in evaluating transgenic hairy roots because of the non-specific feeding of these worms.

[000115] *M. javanica*: Nematode eggs are sterilized using bleach and are used to inoculate hairy roots expressing RNAi. Nematodes are assessed for their growth by measuring knots, egg masses or production of viable eggs. An alternative approach is to microinject dsRNA directly into root feeding sites or into living female nematodes.

[000116] *H. schachtii*: Cultures of this nematode were maintained on sugar beets. Nematodes eggs are sterilized using bleach and used to inoculate hairy roots



expressing RNAi. Nematodes can be assessed for their growth by measuring knots, egg masses or production of viable eggs.

#### Example 5 – Plant Expression Vectors for RNAi

[000117] Modular Binary Construct System (MBCS): An important aspect of the subject disclosure is the Modular Binary Construct System. The MBCS eases the burden of construct development by creating modular pieces of DNA that can be easily added, removed, or replaced with the use of low frequency cutting restriction enzymes (8-base cutters). These constructs are useful for delivery of a variety of genes to plant cells and is not limited to the delivery of RNAi genes. To develop this system, a series of six, 8-base cutter restriction enzyme sites was placed between the left and right Ti borders of a previously created  $\text{kan}^R/\text{tet}^R$  binary plasmid (Figure 1). The production of both  $\text{kan}^R$  and  $\text{tet}^R$  MCBS aids the testing of constructs using different strains of *Agrobacterium rhizogenes* in different plant species. In addition to the MBCS, a series of shuttle vectors were created that aid in the cloning of useful DNA fragments by containing the multi-cloning site (MCS) of a modified Bluescript plasmid flanked by 8-base restriction sites (Figure 2). With six 8-base cutter sites, each site is, preferably, reserved for a particular function (Figures 3 and 4). Because of the close proximity of the *Pme* I and *Sgf*I sites to the left and right border of the binary vector, these sites are, preferably, reserved for gene tagging and enhancer trap experiments. The *Not* I site is, preferably, reserved for plant selectable markers (Figure 5). The *Pac* I site is reserved, preferably, for Plant Scorable Markers (Figure 6). The *Asc* I site is, preferably, reserved for RNAi experiments (Figures 7 and 8), while the *Sbf*I site is, preferably, reserved for anti-nematode proteins. The restriction sites that are denoted in the Figures are, preferably, reserved for the denoted insertions; however, the MCBS binary and shuttle vectors do not require the restriction sites to contain these suggested inserts.

[000118] Plant Selectable Markers for MBCS: To further develop the MBCS, a series of plant selectable markers were added to the MBCS (Figure 5). Plant selectable markers that were added to the MBCS include: pNOS/NPT-II/tNOS ( $\text{kan}^R$ ), pNOS/Bar/tNOS ( $\text{basta}^R$  for dicots), pUBI/Intron-Bar/tNOS ( $\text{basta}^R$  for monocots), and pUBI/Intron-PMI/tNOS (mannitol isomerase $^R$ ).

[000119] Reporter Genes for MBCS: Four exemplary reporter genes are used in the MBCS are provided in Figure 6 and Appendix 2. GUS, a nuclear localized GUS, GEP, and the anthocyanin transcriptional activator *pap/C* genes into the MBCS.

[000120] Promoters for MBCS: We cloned several useful constitutive and nematode-inducible promoters (Figures 6, 7 and Appendix 2). Constitutive promoters include the SuperUbiquitin promoter from pine (pSU) and two promoter regions from the Strawberry Banding Vein virus (pSBV<sub>1</sub> and pSBV<sub>2</sub>). Seven nematode-inducible promoters from *Arabidopsis* were also been cloned.

[000121] The following Scorable marker clones have been constructed and placed in the MBCS, NPT-II binary vector (pNOS/NPT-II/tNOS):

Intron/GUS/tNOS	Intron/NLS-GUS/tNOS	Intron/GFP/tNOS
pSU/Intron/GUS/tNOS	pSU/Intron/NLS-GUS/tNOS	pSU/Intron/GFP/tNOS
pSBV <sub>1</sub> /Intron/GUS/tNOS	pSBV <sub>1</sub> /Intron/NLS-GUS/tNOS	pSBV <sub>1</sub> /Intron/GFP/tNOS
pSBV <sub>2</sub> /Intron/GUS/tNOS	pSBV <sub>2</sub> /Intron/NLS-GUS/tNOS	pSBV <sub>2</sub> /Intron/GFP/tNOS
pKT/Intron/GFP/tNOS		
pKA/Intron/GFP/tNOS		

#### Example 6 – Control of Plant parasitic nematodes using RNAi in planta

[000122] Production of RNAi Vector. The RNAi shuttle vector to be used is adapted from the Modular Binary Construct System (MBCS - See Example 5). RNAi shuttle vectors preferably comprise a promoter, intron, antisense RNAi, stuffer fragment, sense RNAi, and terminator (See Figures 7 and 8 and Appendix 2 for more details). The plant promoter can be constitutive, tissue-specific or nematode-inducible. The intron is necessary to eliminate expression in *Agrobacterium*.

[000123] The anti-sense and sense RNAi molecules comprise nematode-specific sequences and are disclosed herein. These genes are associated with pathogenesis, growth, or other cellular function in nematodes. An exemplary group of RNAi sequences for use in plant/nematode control may be based upon:

[000124] 1. Genes specific for nematode esophageal gland cells.

[000125] 2. Genes specific for plant parasitic nematodes but not other free living nematodes.

- [000126] 3. Genes common to all plant parasitic nematodes.
- [000127] 4. Genes common to all nematodes (nematode-specific).
- [000128] 5. Genes specific for important tissues or cell types.
- [000129] 6. Genes from large gene families.
- [000130] 7. Genes involved in nematode signal transduction or other cellular pathways.

[000131] Appropriate RNAi constructs allow for the formation of dsRNA molecules (the sense and antisense strands join to form the dsRNA). The terminator sequence adds a poly-A tail for transcriptional termination. The RNAi shuttle vector can then be subcloned into the MBCS and transformed into *Agrobacterium rhizogenes*.

[000132] Plant Transformation with RNAi Vectors. An exemplary transformation system for generating hairy roots using *Agrobacterium rhizogenes* is provided below. The RNAi vector once introduced into the MBCS can subsequently (as a binary vector) be transformed in *A. rhizogenes* using, for example, the electroporation protocol of Example 2. Once the *A. rhizogenes* is confirmed to contain the plasmid, it is then used in generating hairy roots (See Example 3). Using this protocol transgenic hairy roots expressing RNAi are isolated, cultured and tested.

[000133] Testing of RNAi Vector for Nematode or Plant Pathogen Resistance. RNAi expressing hairy roots can be inoculated with sterilized nematodes. Infested hairy roots can be observed and the effect on nematodes determined. An alternative approach involves the microinjection of RNAi directly into root feeding sites (giant-cells for root-knot nematode, and syncytia for cyst nematodes) or into living female nematodes.

#### Example 7 – Insertion of Genes Into Plants

[000134] One aspect of the subject invention is the transformation of plants with genes encoding proteins of the present invention. Transformation of plants as described herein can be used to improve the resistance of these plants to attack by the target pest.

[000135] Genes, polynucleotides, and/or RNAi molecules as disclosed or suggested herein can be inserted into plant cells using a variety of techniques which are

well known in the art. For example, a large number of cloning vectors, for example, pBR322, pUC series, M13mp series, pACYC184, pMON, *etc.*, are available for preparation for the insertion of foreign genes into higher plants via injection, biolistics (microparticle bombardment), *Agrobacterium tumefaciens*, or *Agrobacterium rhizogenes*-mediated transformation, or electroporation as well as other possible methods. Once the inserted DNA has been integrated into the genome, the genetically modified-cell(s) can be screened via a vector carried-selectable marker that confers on the transformed plant cells resistance to a biocide or an antibiotic, such as kanamycin, G418, bleomycin, hygromycin, chloramphenicol, or bialophos, *inter alia*. The transformed cell will be regenerated into a morphologically normal plant. The transgene(s) in the transgenic plant is relatively stable and can be inherited by progeny plants.

[000136] If a transformation event involves a germ line cell, then the inserted DNA an corresponding phenotypic trait(s) will be transmitted to progeny plants. Such plants can be grown in the normal manner and crossed with plants that have the same transformed hereditary factors or other hereditary factors. The resulting hybrid individuals have the corresponding phenotypic properties.

[000137] It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application.

We claim:

1. An RNAi molecule, optionally comprising a linker, wherein at least one strand of said RNAi is encoded by a DNA sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 139.

2. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
1.

3. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
2.

4. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
3.

5. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
4.

6. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
5.

7. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
6.

8. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
7.

9. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
8.

10. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
9.

10. 11. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
12.
11. 12. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
12.
12. 13. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
12.
13. 14. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
13.
14. 15. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
14.
15. 16. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
15.
16. 17. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
16.
17. 18. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
17.
18. 19. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
18.
19. 20. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
19.
20. 21. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
20.

21. 22. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
21.
22. 23. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
22.
23. 24. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
23.
24. 25. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
24.
25. 26. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
25.
26. 27. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
26.
27. 28. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
27.
28. 29. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
28.
29. 30. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
29.
30. 31. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
30.
31. 32. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
31.

34

32. 33. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
33. 34. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
34. 35. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
35. 36. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
36. 37. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
37. 38. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
38. 39. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
39. 40. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
40. 41. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
41. 42. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
42. 43. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:



43. 44. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
44. 45. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
45. 46. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
46. 47. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
47. 48. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
48. 49. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
49. 50. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
50. 51. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
51. 52. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
52. 53. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
53. 54. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

54. 55. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
54.
55. 56. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
55.
56. 57. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
56.
57. 58. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
57.
58. 59. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
58.
59. 60. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
59.
60. 61. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
60.
61. 62. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
61.
62. 63. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
62.
63. 64. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
63.
64. 65. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
64.

65. 66. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 67.
66. 67. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 68.
67. 68. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 69.
68. 69. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 70.
69. 70. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 71.
70. 71. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 72.
71. 72. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 73.
72. 73. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 74.
73. 74. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 75.
74. 75. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 76.
75. 76. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

76. 77. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
77. 78. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
78. 79. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
79. 80. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
80. 81. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
81. 82. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
82. 83. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
83. 84. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
84. 85. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
85. 86. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
86. 87. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

87. 88. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
88. 89. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
89. 90. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
90. 91. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
91. 92. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
92. 93. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
93. 94. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
94. 95. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
95. 96. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
96. 97. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
97. 98. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

99. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 98.
100. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 99.
101. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 100.
102. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 101.
103. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 102.
104. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 103.
105. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 104.
106. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 105.
107. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 106.
108. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 107.
109. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 108.

110. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 109.

111. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 110.

112. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 111.

113. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 112.

114. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 113.

115. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 114.

116. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 115.

117. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 116.

118. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 117.

119. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 118.

120. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 119.

121. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 120.

122. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 121.

123. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 122.

124. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 123.

125. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 124.

126. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 125.

127. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 126.

128. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 127.

129. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 128.

130. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 129.

131. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 130.



132. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 131.

133. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 132.

134. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 133.

135. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 134.

136. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 135.

137. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 136.

138. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 137.

139. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 138.

140. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 139.

141. A transgenic plant or transgenic plant tissue comprising an RNAi molecule according to any of the preceding claims.

142. A method of disrupting cellular processes in a nematode comprising the steps of:
- (a) providing a composition comprising a compound according to any of the preceding claims; and
  - (b) contacting a nematode with said composition.

143. An isolated promoter comprising the following nucleotide sequence:

aacagcccaagataaacagaaaagtcaaaggtgttcgaaa  
gaccacttgtgactaaggatcatttcacccataattatctggtagca  
cagactcatgataaetgcgaggaacacaagttctttacagtcgattc  
aaagacactttctctttacggtttcattgaaggagccgaccagaat  
atgtcagagaagcttttctactgtgggttaatttcattaatctatcca  
gggtgaaaacctcaaggagatctctctctcccaaaagacctctacag  
ggcaatcaaaaactacagaaccagagtttgtagtgcacagagtagac  
caatctacctgagaatcacgagtaccttcctagagtgggaaaatgat  
gacatccttattccataccactggattgaggtaggactatccaatgg  
aaaaattccatgggacaagtcataaagaagaccgcaacagtcgagt  
atcttccagagataaactgcactcagacctaaaaggataaaaagcagta  
tataatcagtgtactaagatcttcgcagattcaaagaagaagcttaa  
ctatgctgatgacaagataattctaataagcaattattcagaattaa  
tcaaggagaaagaattaataactctttcagaatatgaagcccgttt  
acaagtggccagctagctatcactgaaaagacagcaagacaatggtg  
tctcgatgcaccagaaccacatctttgcagcagatgtgaagcagcca  
gagtgggtccacaagacgcactcagaaaaggcatcttctaccgacaca  
gaaaaagacaaccacagctcatcatccaacatgtagactgtcgttat  
gcgtcggctgaagataagactgaccccgaggccagcactaaagaagaa  
ataatgcaagtggtcctagctccactttagctttaataattatgttt  
cattattattctctgcttttgctctctatataaagagcttgatattt  
catttgaaggcagaggcgaacacacacagaacctccctgcttaca  
aacatgtattgtagctaaacctcttaggag .

144. An isolated promoter comprising the following nucleotide sequence:

tggtggggacaatggatccggtctgcgtagcaacaaggctg  
aaaaagattaaacagaaacctgtgatcattagcgttggaccaccacc  
aaaacctcctgagccaccaaagcctccagagcctgaaaaaccaaagc  
ctccaccagcacctgaaccaccaaagcatgtatgcaagccaccttac  
tgcaacagttgtgatgttgtgtctgttactacctatgaaagtggaag  
cggctgcaccattctttgagtcataatcgcgtagcatagccttcac  
gttaagtccctgtatcttagccaataactaattcatcatgttctcatgct  
tttttggtttatttctttttctcaaatatgaatctctgttgtttgtcc  
ctccccctgtttataattagtcgcttctttgacacaagaagtctcatg  
agttcatgctaaagaaaataaaagttcaaattaaaacaccaaagtgt  
tgattaatctccataaacctgtgaagcagaaagttagtcattgttgac  
ctgaacagagccttaggaagtccttgaaggacatatcttcaagtgtta  
ttgggtcgtagcactcttagggccattacttcattgagcccattaa  
attatgcaaaacaagaaatgagacatatggaaacattagggttctta  
caggaaaaaataggaaaaagcagggacaactaaacaaaaattcagaa  
acaagaggcaagtggacgaccacggcgtaagatcaacatgtggtgat  
gtgcatgagaccaagaccattttttctcgttcttcaacgcacacttg  
gtctttttcttatgtttgttgcatcttctttattaggcagaccctctct  
cttttttaatataggatagtaaaaaatatatgattttattttgttgaaa  
cattttgagttaaaacctaaacttatagtaagcatttgtagagtga  
tttctctatcagacatctatcaacatgacctctaaccccccaaatatt  
gatgaaactactttaagtagtaaaacctaaagcaattaaaatttctct  
ttaaattagtagtttgtgtaaattaatgacatgattgctgcgaaag  
aaatcaaaacagttatatcggtgaacttaggagaatgttttatatcgt  
gtttcaacacatgattgctagcatatgtgtagggtgctgtagacgtta  
cataacaatcatcactcgtaaatatcaaagtgggtttctgagagaaac  
aaagggttatgattttcccaactgcactagttgtgtattgtttcttt  
cacacgtatgcttctgagttctgcccagggtggaattaaagcagag  
ttgggagagatcataatttattagggttcgttatgctcaagtcatga  
cgtaaaatgaaaatttgtttttattctttcaccaacacaaaagaatag  
ctagttatctctttttttatatataacaattcatgaagttgatcagc  
tttatacacatcatccaatcgaattgctaatttagagatggaaatat  
caggatagagccaataagatatcaaatccaatggaccattttctcc  
atgtgctaattcatacaatctgtttttgtctgctttatttgatgatg  
atgctgagcgtttttaagtgtgaactaagatctagctaaccaaaaca  
aagatgggtctctctgtctttgtcgtataagagcaagagagtggttt  
gattcaatttttaaaattctaaataaaactccaaccgtgaatccagc  
catgaaactctttttagaaaatccttttttataacaaataattcctc  
tgcctctctctctctctctgtttatttcaccttttttggtttctttag  
ctcagaaaaagcccatctttttttctattcttggtttattttaatca  
tactgtgcgtttctacaaagttgttcctttctcttcaactctctc  
actcacagtcacagagatctgtttctttttttttttgttttcaactc  
ttctcttccagt.

145. An isolated promoter comprising the following nucleotide sequence:

agcaaagcaagaacaccagagaagaagaaaagcactacaga  
gaaaaatgtgagcttaagcgtctccaacaacacttctctgggagtc  
taaaggatgctgcaaaaagccttgggtggtgagacttccgcatatttc  
caagcatgggtttatttttgttagcacacaaactatctgaccctcga  
cttggattttcttctgcagtttgtccaactacattgaaacggatatg  
caggcaacatgggatcatgaggtggccatctcgtaagattaacaaag  
tgaacaggtcactaaggaaaatacagacgggtactggactcgggtccaa  
gggtgtagaaggaggactaaagttcgactcagcaactggcgaattcat  
tgcagttagaccttttattcaagaaattgatacccaaaaggtctgt  
cgtctcttgataatgatgcacatgcaagaagaagtccaggaggatatg  
cctgacgatacttcattcaagctccaggaagctaaatctgtcgacaa  
tgccattaagttagaggaggatataccatgaatcaagcaagaccag  
gtaagaacttctctatccataaaccatagatggagcgttagaatct  
taatccattttcagtttttgcaggatcattcatggagggttaatgcta  
gtggtcagccatgggtctggatggccaaagagtcgtggttgaatggc  
agtgaaggaataaagagcgtttgcaacttaagctctgtggaaatttc  
agatgggaatggatccaacaatccgatgcagtggtcagttgttgaa  
ctaaccaatccatgtcatgcagcatatcagattcatcaaatgggtca  
ggcgcagttctgcgtggaagctcatctacttccatggaagattggaa  
ccaaatgagaaccacaacagtaatagcagcgagagtggtcaacaa  
cgctgatcgtaaaggccagttatagagaagacactgtacgtttcaag  
ttcgagccatcagttgggtgtcctcagctctacaaagaagtggaaa  
acgttttaactgcaggacgggtcggttcagctgaagtacttggatg  
atgaagaagaatgggtgatgctggttacagattctgatctccaagaa  
tgtttggagataattacatgggtatgggaaaacactcgggtgaagtttct  
cgttcgtgatttgtctgccccctctaggtagttctgggtggcagtaatg  
gttatcttggaaacaggcttatgacgtcgtaagacatagacacacaca  
gttatgtattcccagtgaaagaatgttgtttatttctctagatatta  
gtatgcttataaataggcatgaaggagaaagacaattttgggtatagt  
ggagtccagcagaaaatgtatatgtttttcgttttatatgaatcag  
agaataaaaagttggatgttatatctacgttgctaattgttgtaacctgc  
tcaccatctttcatataagaaaagagaacacttttagttatccctg  
tgatgcagaatcgtattctttgttatctctccattcctgtggaaacc  
aaciaaagtcactaaatttcgggttaattgggtgggtttttaagtcaa  
cgaggacttgatttttagttgggtctgggcctataattgtgttcatca  
ttgggttttttcccccttatcagtttaacgtccatatccatatcttt  
ttctttttttaacggcaagttcatatccatatcttatgatgtgcct  
aaaagagggagaagatgcgaagacagaattttcatatttgaaagggt  
tcgatatcgatattgggaaacgaatcaagggtcaaaaaactcagttca  
atagttgaaatttaaaaaattttattaattcaatccgattgggttcgt  
tttgttatgggttcgggttctatatcatcaaaccaatcggtttgggtcct  
aaagataattataaatattcaccaacaccagtggttaaacacatatca  
acaaacctaaagttagataaacaagaga.

146. An isolated promoter comprising the following nucleotide sequence:

aattggcactcttcttctgctgggttccaaaagaaacgaat  
caatatgtgcaacaagaagagctccagaagcagtcattcttctaaaat  
cttaatctaacaacagctcaagaagaaaaaattccatagctagaga  
gaacacaaagtcacaagacgacgtcgtagaggcacaagtcacacct  
gaatggcttaagccgaactgagtggttttgactagaccatcatcaga  
aaagtcctccaagacggtagtcggatgttagatcgctcaagtaatttt  
tggttttggttggtctcacgttttcagctgcccatttgatttcagttt  
gggcttttccttatctctaaaggcccaatttcatttaggtttagttt  
atttgatcattatccttactataaaggcttcgcctttcgagaaattt  
agggtttcttctgtctgtctcgtcactcaggtttgtgcctcaacgac  
tgcttcacttctagcttgattcttcttcttcgtttatatgtatactg  
tacattagattattcttgtttctcgagcttctgctatagattttgat  
tcttttttttggttggtcttctgttttcgtttccaggatcagatcttagct  
aaattgagacaagctcaaaatgaggtacttgacgcacatctcttaoatt  
cactgtttaattagagaacaatacgtctctgaatcgtgattcagaga  
cgtattgttcttctgtcatatgcaataagtttaattagagaacaata  
cgtctctgaatcgtgattgttttttggttggtgcgttattgatagctt  
tatgatgttaatagcttaggattgacacgaagtgttctgcagtttt  
gcataaatgctctttactaaggcctctaaatttggttgacaaatcta  
aatcttgccctataaaaaatttaggtgtattaaagataagattattttg  
tatggtagtgctctataatgtgggttggttcattgttgagggttgtaacg  
ttgtgtattttttgtttgttttagttaatttgcttaactctgttctttg  
tgggttaatacagtaagcttcagagtgaggccgttcgtgaagccatc  
actactatcacagggaatccgaggcaagaaacgtaactttgtcga  
gactattgagctccagatcgggtctgaagaactatgaccctcaaaagg  
acaagcgtttcagtggtatctgtcaagttaccacatatcccccgctcct  
aaaatgaagatctgcatgctcggagatgccagcatgttgaagaggt  
gatatatcttttcatggaaattgatcattttgtgctctgtttcttgt  
ataatgggttttggtgctcattttcatttggtggctctattagtttcatt  
tgatgttgatatgtcttctgaatgtagatgcatgatgttttcggaa  
tttggtcattgtttatttaggttctatttcttgcataattaaatatt  
tgcttattttcatcttgtatcttttcgtaggctgagaagatgggggtg  
gaaaacatggatgttgagtccttaaaaaagcttaacaagaacaagaa  
actcgtcaagaagcttgcaagaaataccatgctttcttggtcctctg  
agtctgtcattaagcagattcctcgtcttcttggtcctggctttaac  
aaggcaggcaagttctggctacagctaataatccattgttcttcttt  
acatccgttttgatttttggtataggttttagtagtctatttcttttgt  
caatgtctttttgatacaatgccaatcctttatcctgtgagattatg  
cttctttgatgattcttaagtaacattcctttgctttactttacaca  
ggaaaattcccaactcttgtagaccaccaggaatccttgaggtcaaa  
ggtgaatgaaacaaaggcaacagtgagttccagctgaagaagggttc  
tgtgcatgggagttgcagttggtaaccttt.

147. An isolated promoter comprising the following nucleotide sequence:

```
ttggcaaactgagatataagaggggaaggtgattttcatgcaa  
atTTTTTTTTTattTTTTTTTgaatgaatgcaaaatttattcaaaaa  
aaaaaaacctgggctacatcaagtacttcatttctgagtttttgaaa  
aatctaaagacaacaaaagactttacaatttaataaaaaaataataa  
aaatactttatcactctcaacgaaattgttgatttaataacgtatct  
cttggtaaaacagcgttttatttgacgaaattgttataaatgaataa  
aatgataatagaaactagtgtggtacgtaaaatacctctcatttggc  
aaaataacggttatgtatcatgagttatgcatacgacagcgtgctta  
aatagtgtgctttcaggagaaaatatataccaagttatttgctgaaa  
ttaccacgcaaactcgagggtcgaatggcaaaaataaaaaaccaatgt  
catttccttaatgtattaaggtcatttaataaaaattgtacactttt  
ttcacctgtaagcgttccaaagtgtagaatggataactagaaggggtc  
aaaggtataatattaataagcgaactcactttttgcccaagtgattt  
cacttcttacattttgcttgatatagttacccaaaagtgtatatatat  
tccttatatacaattgttctattttctggattataaggggaataagaa  
aaaagaaaagagagagtataataataacttttataaagtgatgtta  
gattctaatttgtaacgaaaagttcaaagtgaagaaaaaacgaaaa  
agtttttctgttttgttttatatctatagccaagaaagtttctcaga  
tttacaagaagttaactgagaaaaacaaaaaaaaaacttatgaagca  
tgaaagactaattaacgaggtgattaattttgagacaaattaaacat  
cgaattaaaagtaacatttgagggtttatatgttatatatgtgaca  
tgataagtccgattcatgactaatgtatatctggaatctaacatgga  
agaatagagaacgaagcagagccaaggtcaacttgccagacacgaat  
caacagatttgtgaatgagaccaaatacaatgggtcataaacgggtggg  
tttaaacgggcaagtcaccttggtcgaattccattcggtatttcctt  
catgcaagaccctctgatataccaaagactccattacaatattctt  
ttcgatcacgagctacttattttcaaagtgtgttacctcttctgtgac  
tcttgtgttggtggttaaagcctagtcgagatgtgtcggtatatata  
ggcatatatacaaatgcgacaaaataagttatattatattgtttaa  
tttctatattccatttctatatgcatggctgggatttttgacaaaa  
ccctaattcaagaatagaatccaaaagatgggatcaaagaatataat  
ctaattgggtgaccacattttccgatttaattcgcatagttaatatt  
ctttccactactttatgccgcagaaatttgtaattaaagtaagacaaa  
gaaatacagatataagatgggtcgtagaaaccagtagaggaatttcat  
ttttcggtggataagtgggaatttaataagagaatgggtctttactctt  
tacagtgggaaatgggaatagtagccattataatttcatcagattc  
tatatatgcatgtttgtataagctaaaaataaaatacgtttaagcattc  
ttcaaaaaaattttacaagttctagagactctcttaacgtcggcaatt  
tatattctactttacatgacactttcaggaaaagaaaactatactca  
ctagcagatcattaaattttctttttctttttttgaatgaaccttag  
ttgtgggtttttatttttggtagctagaaacttcagtgtttttttcc  
gccaatggtagtgctttgatgatgggtccgg.
```

148. An isolated promoter comprising the following nucleotide sequence:

```
caatcaaggtaacgaaggaggatcagcgaaaggatgggcta
tatttggagtttttctcgtgtaagtaatgctttgtgatcttcca
tgcgacatataactgaagaataaactcaactcattgtgttctgggtg
tggttcttctgatcagattcctcgttgcatctgcacttttctgctgt
gggggctttatttataaaacaagagtagagcgtgtggtaatcttcat
atcttctacaattccacttccattctctaattattctctcacgtga
tatacacacactcaatcactgatgtactcgtatggatgcagcgtgga
actgatgcattgccggggatgtcacttctatcgggcttactagaaac
tgtaagtatatacaagaaaactcaaaaggattccatttatgcaaaatc
taagagaaaagctcactgtggtctttgggtacaatttatggatctctc
aagagacaaatgctatgtaagctaattgatttttgggtcttgataaaca
ggtagagtgaagtggaacaaagctactcaagaactgaagacatcaaca
atgcttttgccaatgaagtctcatgggaccgctcttccgcatcttct
actcaagcgacaacaacacagagaccaagtgaagaacatatgggtgc
gatctaattttgtcaagtgcctcacaagaggtagtgttcaagccat
ggtagtggaacgcttgtgatctgcgatttctggattttgctttgtatg
tttattttctaccttctagaaagagggtcaaaaagttaatagcttcac
cgtgagaatgttggttttaccagattcatgtgctatgatagaaaag
acaaagcaaaacaagagttcttcttcttgcttaggttacaagaacaaga
gtatcggtataaaagtcaacaaagattgaaacatattttgtcaagg
agtgggttagaatctcttctactctcttgcccttctcactaagacaa
aaaaaagacttggactttgtctaagggtttgtggatattattaacca
agtccttttgcaaaaagtaatatgttttttcgcattcctcttttag
aatttagtttaattctaggctttatatgtggttattactttcttgaaaa
atgatctgtttattctattcatacttgggttacctcgctttttatctt
acttctacaaaaggattatcagtgaaagttagtctcttactctcacc
ttccgaaaataaaaacaaaatatcgatacttctagatcaaaccaagt
tgattaaaacatccctattccctacgattctgatcttgagatatatt
atcatgttaagatctaaattgacaagaaaactgatttttcatattcta
gtaggaaaaataattactattagtgatcatgattgtcgaccgtaaga
ggtaggttttagttactctccatctttcttgaagaagtcagaaagtca
gaaattatatcaaatcaaacatcaatattgaacacatatatctgtat
ggtttttatgttttagaaaaattccaatatattatatattcctagggaaaa
agaagcttattcttcaaattattgttatgagtcgttaaaatatggat
aaaaatataaagtctaaatatataaaactcagtttgctttgctttta
cctctccaagtcctcaaagtc aaattaatttttagttaattaaaccaa
aaaagggtttattagtc aaacttagcatgcaatgctgggtaccaaacc
caagcattagtcctcttttaattctcttttctccaataagtttttac
aatttttaattgtttgcatttcccttgattatttatcttcatcccaa
tttagctaataccaactccgtttcttattcttccaagtccttttcta
taaatacgttcttcttccctcttatttcatatcactcaccacaaag
tcttctcatttccctcat .
```

149. An isolated promoter comprising the following nucleotide sequence:

atgttgtagtgagtgaggagaagaagaggggaaacaaaggtatt  
tatttgtagcgagttttgttttgtagcggtttgtctgtgttcaa  
tgttgacgaaacgagtgagagagtgctgattattaaagaaaaccct  
aattaagtcagaccccgccggtataaaaaatagtcaaaaagtaggaaa  
acgcgtgtgtgagtgagacagagacagccattgtttgctttatggg  
cttataagcgagacgtgttaattgggcttttctttatggccgaaa  
acaaaaagaaacgtcgctgagagattcgaaactctcgcgggcagagcc  
catgtacttagcaggcacacgccttaaccactcgccaaagcgactt  
gttgctatgagttagacaaaaatcattaaaattctctattatgatttc  
tcatagtggtgtgtatattgtggatctactaaaaattctttgttat  
tattactttattttgtgaattagtttgatataggtaagtacaaagtt  
aactttattttactcaaaatttatcagattaactgattttatatt  
gtttccttttggtatatagacgtactatagtttttagaaaaaccataa  
gattcctttatatttcatagagtgagagatgagatgagatcttggc  
tgagagaagaaataagttccacgaggaggactcttttttttggtga  
agacgaggaggaggactcttgggtgatccagtctttacgttagacat  
cgccccctacattttatttgctttctctatcaacatggcaggtaaaa  
atcttcattcaaccgaaccaaccaaagtctcttcccaataatattca  
agcaccatcctttgggaaactcatacactacagtctacactcttt  
cattttctttcaacgctcaacttaacaaatgatatagtctagttgtc  
aattatatgttttaattagtgttttcacatcaaattctgggttgata  
tttgatgactattttcggaacatctcaatgtcccgcaaatacaatc  
gtctatcatatataatcccgtagcttgattcttatagatagaataa  
tatggcgtgatctttataatataacatatagaatcgtgtagatttat  
ttattttatttttatatcgcatataattgcaaaatacttatatat  
gtttgttatatatgataccattttatagttacttaaaaaaagttaa  
gcgataatatatatatacaactttttataacaaaaaagtataaacac  
atggtaaagaaaaataaaaaatgaagacatgggtgtgacacgaaaatgg  
cactaaatatacatatataatagatagctacaatatcccatcataca  
cacttttttaattgactaatacataacttacacacttttttaattga  
ctaattcataactttttatcattgtcaacatgcaaattcatatttcc  
gttgaactattattcttattttgtttttaaaagaagggttctctggg  
aataaaaaatagatttccaaatgacgttagagcaaaaaaaaaaag  
gttgctctgggtctggtaaaatgaaaaagcaaaagcgtcttgggtatagaa  
aagtaataatactgcctcctaatttcttcgtccttctaccgaagaatc  
tctccactcttgccctctttcgaaaccctaaaccagaagcaccagat  
tttttcaactttttccagagaacaatagaaaacccaacttggtgtc  
tctaggggtttctttattccttctcatctttggattttcttgggtca  
tcattttggaagcttaccaccagcgaaaaaattataacttccatcg  
attcctggcttctctctctcgctctctctgcatgtgctaaatcgccg  
gactgatectcactgtcacctctgtt .



150. An isolated promoter comprising the following nucleotide sequence:

gattaggggtttgagttgtcactggaaagaggtttgattgt  
gagtgatgatggagagattatgaaggagtttgtgtgtatttatagag  
gagttaggggtttgaggtttgatgagaagtaggtttgaagaagtttt  
gttgttgcaacttatttagagttacttgttccacaaccacaagtaag  
attggtcacttctaagttctaactagaaacaacccatgacacatggag  
atctcagctaacctagtttaattgtatatgtatttatattttatttaaa  
tattataaaaataaaaataaattttcacaaataaaaagaactacaaaaaa  
gtgagaaaaataatttgataaacaatttagaaaattagtatatcaa  
taaataaatttataatccgatgggttttgccttttgggtttggcctttg  
tttgaacttcgatgagtgactatgtatagcgaaaacaattcggtttg  
tttttgggttaattttaaaaaatacaagcgacaatatctgatgagaa  
taggtgaaaagcaaataatatcagtttaattggaaatatttactttt  
ttacaactaatattttgtttgggtcaaccaacaaatagatttaattaa  
ttatgggttatgagctttttatttggttgacagtatatatatgttaa  
aatagtgatattgcatggcggaagggtccggaagcaacacatatctcc  
tttttaatttttttttaacaagaataacatgttaatttttttttgga  
aatttaataaagaatacatatttctaatttttgcgtcagatagatgat  
taaagagtggtgtgtgtttttttaacaaacaaggaatacattatacata  
tttcatatttctctcgacattgtttgtttttttaaaaaatagattaa  
agagtcctacgaagctaagtagctaacgaagacttgaaatgagaagaa  
gacgagaatcttttaatatattttttgttaagcgataatatatttgaaaa  
ttaataaatatagattaaaggaaataacaataacgcagatatcggtaa  
gtcatagaaaaaaagaaacaacacaaacttacataaacatgtttcct  
aatttgtaatggagtaaaattccttcttttttttttttttttgattt  
ggattccaattagtaaagaactcaatgactataaataacctttaacc  
ctctcattatttcttactatcaattgattaagctctcgttcctaaga  
aagcaatagacgaacaagaacccatcgaagaacacaaatctctcttt  
gaagttgtcgataatgttagtacaccgttacttcgtccaagactttt  
ttgccgttcggtttcttacaaaacaaggatttggttaccattacttt  
tgtcgtaactcctttttacatgtacgtcaaaaagtggttcctcgctc  
cggcttgaagaaacgaccttcttaccacaaaaagcttattttaaac  
cgtctaaaaccggaaaatctcaatctaaaccggatacgggttcatgag  
aaaccgattcaaacaccgagtgagaagtagaattttttgatgggttc  
cgtcacaatgtgtgctgctccttcgccaagacatgtaccgattccga  
tattttgtgggtgtaaagatgatcaaagagtcctcaaagctaagcacg  
acttgaatgagaagaagaagaccaattactcaattagattttgtttt  
gtggagcaattattgtctatttatctttgttttttagcaaataatctg  
tatccactaatcttcacagtacttgactaacaagaagtaaagagttt  
tcttatttccaattgttttttaattctgatacttttttcataatttta  
caatgtttgatgaaaaaaacattcaaacctaaattttctttttttg  
gtatgaattcaaacctgaattacttttgacgaggacccgacggtata  
aatagggatctcccaacaaacaaaaagggt.

151. A transgenic plant or transgenic plant tissue comprising an isolated promoter according to any of claims 143 through 150.

**APPENDIX 1**

<b>SEQ ID NO:</b>	<b>INTERNAL IDENTIFIER</b>	<b>FUNCTION OF POLYNUCLEOTIDE / GENE</b>
1, 2, 3	2293133	glyceraldehyde-3-phosphate-dehydrogenase
4, 5, 6, 7	7143495	Histone H4
8 & 9	7143515	ATP dependent RNA helicase, mRNA sequence
10, 11, 12, 13	7143527	nematode specific
14 & 15	7143602	protein serine-threonine phosphatase 1, catalytic subunit
16 & 17	7143612	40S ribosomal protein S4
18	7143666	cytochrome p450
19, 20, 21, 22	7143675	Neuroendocrine protein 7B2
23, 24, 25	7143839	nematode specific
26	7143863	40S ribosomal protein S17
27 & 28	7144016	vacuolar ATP synthase subunit G
29	7144025	malate dehydrogenase
30 & 31	7144060	J2 pcDNAII Globodera rostochiensis cDNA similar to Bystin, mRNA sequence
32 & 33	7144225	similar to arginine kinase
34	7144354	pyrroline-5-carboxylate reductase

SEQ ID NO:	APPENDIX 1 (cont.) INTERNAL IDENTIFIER	FUNCTION OF POLYNUCLEOTID E / GENE
35, 36, 37, 38	C10	ribosomal protein L18a
39, 40, 41, 42, 43	C118	ribosomal protein S11
44 & 45	C122	ribosomal protein L16/L10E
46 & 47	C127	FMRFamide-related neuropeptide precursor
48	C129	ADP-ribosylation factor 1
49	C130	ribosomal protein L11
50	C137	nematode specific; conserved in <i>C.elegans</i>
51 & 52	C138	ribosomal protein L7
53	C145	ADP/ATP translocase
54 & 55	C148	troponin
56 & 57	C154	calponin
58	C16	translation elongation factor EF1A
59 & 60	C18	40S ribosomal protein S16
61	C27	ubiquitin
62 & 63	C46	nematode specific
64, 65, 66	C48	ribosomal protein S3AE
67	C59	40S ribosomal protein S5/S7

SEQ ID NO:	APPENDIX 1 (cont.) INTERNAL IDENTIFIER	FUNCTION OF POLYNUCLEOTIDE / GENE
68	C8	glyceraldehyde 3-phosphate dehydrogenase
69 & 70	C82	60S ribosomal protein L30/L7E
71	C90	glyceraldehyde 3-phosphate dehydrogenase
72	C135	nematode specific
73 & 74	C206	predicted troponin
75	C227	cytochrome P450
76	C238	vacuolar ATP synthase subunit G
77	C246	40S ribosomal protein S4
78	C308	FMRFamide-like neuropeptide precursor
79	C342	ubiquitin
80 & 81	C344	nematode specific; conserved in <i>C.elegans</i>
82, 83, 84, 85	C370	40S ribosomal protein S5/S7
86	C426	nematode specific
87	C458	histone H4
88 & 89	C481	ribosomal protein L30E
90 & 91	C556	nematode specific; conserved in <i>C.elegans</i>

SEQ ID NO:	APPENDIX 1 (cont.) INTERNAL IDENTIFIER	FUNCTION OF POLYNUCLEOTID E / GENE
92	C628	ribosomal protein S17E
93 & 94	C665	malate dehydrogenase
95 & 96	C669	malate dehydrogenase
97	C694	ribosomal protein S3AE
98 & 99	C709	ADP/ATP translocase
100 & 101	C714	ADP-ribosylation factor 1
102	C721	calponin
103 & 104	C726	ribosomal protein L11
105	C736	nematode specific
106 & 107	C773	troponin
108	C834	nematode specific
109	C860	bystin
110 & 111	C863	troponin
112 & 113	C883	translation elongation factor eEF-1A
116	C888	40S ribosomal protein S16
117	C898	glyceraldehyde 3-phosphate dehydrogenase
118 & 119	C935	peptidyl-glycine alpha-amidating monooxygenase
120 & 121	C937	calponin
122 & 123	C942	peptidyl-glycine alpha-amidating monooxygenase

SEQ ID NO:	APPENDIX 1 (cont.) INTERNAL IDENTIFIER	FUNCTION OF POLYNUCLEOTIDE / GENE
124	C954	arginine kinase
125, 126, 127	C969	calponin
128 & 129	7235653	ribosomal protein L18A
130	8005381	neuroendocrine protein
131	7235496	pyrroline-5-carboxylate reductase
132 & 133	7275710	protein phosphatase pp1-beta catalytic subunit
134	7923685	nematode specific
135	7641370	40S ribosomal protein S11
136 & 137	7923404	nematode specific
138	7797811	ATP-dependent RNA helicase
139	7143613	predicted phospholipase D

## Appendix 2:

### Exemplary genes used for RNAi vectors.

#### Promoters:

##### *Constitutive:*

##### **Super Ubiquitin from Pine**

CCCGGGAAAACCCCT CACAAATACATA AAAAAAATTCTT TATTTAATTATC AAACCTCTCCACT ACCTT  
 TCC CACCAACCGTTA CAATCCTGAATG TTGGAAAAAACT AACTACATTGAT ATAAAAAACTA CATT  
 CTT CCTAAATCATAT CAAAATTGTATA AATATATCCACT CAAAGGAGTCTA GAAGATCCACTT GGACA  
 AATTGCCCATAGTTG GAAAGATGTTCA CCAAGTCAACAA GATTATCAATG GAAAAATCCATC TACCA  
 AACTTACTTTCAAGA AAATCCAAGGAT TATAGAGTAAAA AATCTATGTATT ATTAAGTCAAAA AGAAA  
 ACCAAAGTGAAACAA TATTGATGTACA AGTTTGAGAGGA TAAGACATTGGA ATCGTCTAACCA GGAGG  
 CGGAGGAATTCCCTA GACAGTTAAAAG TGGCCGGAATCC CGGTAAAAAGA TTAATAATTTT TGTAG  
 AGGAGTGCTTGAAT CATGTTTTTAT GATGGAATAGA TTCAGCACCATC AAAACATTTCAG GACAC  
 CTAAAATTTTGAAGT TTAACAAAAATA ACTTGGATCTAC AAAAAATCCGTAT CGGATTTTCTCT AAATA  
 TAACTAGAAATTTTCA TAACTTTCAAAG CAACTCTCTCCC TAACCGTAAAAC TTTTCTACTTC ACCGT  
 TAATTACATTCCTTA AGAGTAGATAAA GAAATAAAGTAA ATAAAAGTATTC ACAAAACCAACAA TTTAT  
 TTCTTTTATTTACTT AAAAAAACAAAA AGTTTATTTATT TTACTTAAATGG CATAATGACATA TCGGA  
 GATCCCTCGAACGAG AATCTTTTATCT CCTGTTTTTGT ATTAAAAAGTAA TTTATTGTGGGG TCCAC  
 GCGGAGTTGGAATCC TACAGACGCGCT TTACATACGTCT CGAGAAGCGTGA CGGATGTGCGAC CGGAT  
 GACCCTGTATAACCC ACCGACACAGCC AGCGCACAGTAT ACACGTGTCAAT TCTCTATTGGAA AATGT  
 CGTTGTATCCCCG TGGTACGCAACC ACCGATGGTGAC AGGTCTGTCTGT GTCTGTCTGCGT AGCGG  
 GAGAAGGTCCTCAT CAACGCTATTAA ATACTCGCCTTC ACCGCGTACTT CTCATCTTTTCT CTTGC  
 GTTGATAATCAGTG CGATATTCTCAG AGAGCTTTTCAT TCAACCCGGG

##### **Strawberry Banding Vein Virus 1**

aagctttttcactgtgggttaatttcattaatctatccaggtgaaaacctcaaggaga  
 tctctcttctcccaaaagacctctacagggcaatcaaaaactacagaaccagagttt  
 gtagtgacagagtagaccaatctacctgagaatcacaggtaccttccctagagtggg  
 aaaatgatgacatccttattccataccactggattgaggtaggactatccaatggaa  
 aaattccatgggacaagtcatataagaagaccgcaacagtcgagtatcttcagaga  
 taactgcactcagacctaaaaggataaaagcagtatataatcagtgactaagatct  
 tcgagatttcaaagaagaagctt

##### **Strawberry Banding Vein Virus 2**

Gtttaaacacagcccaagataacagaaaaagtc aaaggtgttcgaaagaccacttgt  
 gactaaggatcatttcatccataattatctggtagcacagactcatgataaactgcga  
 ggaacacaagttctttacagtcgattcaaagacactttctctttacggtttcattga  
 aggagccgacccagaatattgtcagagaagcttttctactgtgggttaatttcattaat  
 ctatccaggtgaaaacctcaaggagatctctcttctcccaaaagacctctacagggc  
 aatcaaaaactacagaaccagagtttgtagtgcacagagtagaccaatctacctgag  
 aatcacgagtagcttccctagagtgaggaaaatgatgacatccttattccataccactg  
 gattgaggtaggactatccaatggaaaaattccatgggacaagtcatataagaagac  
 cgcaacagtcgagtatcttccagagataactgcactcagacctaaaaggataaaagc  
 agtatataatcagtgtagtaagatcttcgcagattcaaagaagaagcttaactatgc  
 tgatgacaagataattctaataagcaattattcagaattaatcaaggagaaagaatt  
 aataactctttcagaatatgaagcccgctttacaagtggccagctagctatcactga  
 aaagacagcaagacaatgggtgtctcgatgcaccagaaccacatctttgcagcagatg  
 tgaagcagccagagtggtccacaagacgcactcagaaaaggcatcttctaccgacac  
 agaaaaagacaaccacagctcatcatccaacatgtagactgtcggtatgcgtcggct  
 gaagataagactgaccccgaggccagcactaaagaagaaataatgcaagtggtcctag  
 ctccacttttagctttaataattatgtttcattattattctctgttttgcctctctat  
 ataaagagcttgatctttcatttgaaggcagaggcgaaacacacacacagaacctccc  
 tgcttacaaccatgtattgtagctaaacctcttaggaggatattc



**Nematode Inducible:****Trypsin Inhibitor from Arabidopsis (clone#6598343)**

cccgggagcaaagcaagaacaccagagaagaagaaaagcactacagagaaaaatgtg  
agcttaagcgctctccaacaacacttctctgggaggtctaaaggatgctgcaaaaagc  
cttggtggtagacttccgcatatttccaagcatgggtttatttttggtagcacaca  
aactatctgaccctcgacttggattttcttctgcagtttgccaactacattgaaac  
ggatatgcaggcaacatgggatcatgaggtggccatctcgtaagattaacaaagtga  
acaggtcactaaggaaaatacagacgggtactggactcgggtccaaggtgtagaaggag  
gactaaagttcgactcagcaactggcgaattcattgcaggttagaccttttatccaag  
aaattgataccaaaagggtctgtcgtctcttgataatgatgcacatgcaagaagaa  
gtcaggaggatatgcctgacgatacttcattcaagctccaggaagctaaatctgtcg  
acaatgccattaagttagaggaggatataccatgaatcaagcaagaccagggtga  
acttctctatccataaaccatagatggagcgattagaatcttaatccattttcagtt  
tttgcaggatcattcatggagggttaatgctagtgggtcagccatgggcttggatggcc  
aaagagctctggcttgaatggcagtgaaaggaataaagagcggttgcaacttaagctct  
gtggaatttcagatggaatggatccaacaatccgatgcagtgagcagttattgttgaa  
cctaaccaatccatgtcatgcagcatatcagattcatcaaatgggtcaggcgagtt  
ctgcgtggaagctcatctacttccatggaagattggaaccaaatagagaaccacaac  
agtaatagcagcgagagtgatcaacaacgctgatcgtaaaggccagttatagagaa  
gacactgtacgtttcaagttcgagccatcagttgggtgtcctcagctctacaaagaa  
gttggaacacgttttaactgcaggacgggtcgtttcagctgaagttacttggatgat  
gaagaagaatgggtgatgctggttacagattctgatctccaagaatggttgagata  
ttacatggtatgggaaaacactcgggtgaagtttctcgttcgtgatttgcctgcccct  
ctaggtagttctgggtggcagtaatgggttatcttggaacaggcttatgacgtcgtaag  
acatagacacacacagttatgtattcccagtgaaagaatggtgtttatttctctaga  
tattagtagtcttataaataggcatgaaggagaaagacaattttgggtatagtgaggat  
tcagcagaaaatgtatatgtttttcgttttatatgaatcagagaataaaagtgtga  
tgttatatctacgttgctaatgttgtaacctgctcaccatctttcatataagaaaag  
agaacacttttagttatccctgtgatgcagaatcgtattctttgttatctctccatt  
cctgtggaaaccaacaaagtcaactaaatttcggtttaattgggtgggttttaagtc  
aacgaggacttgatttttagttgggcttgggcctataattgtgttcattcattgggttt  
tttcccccttatcagtttaacgtccatatccatatctttttcttttttaacggcaa  
agttcatatccatatcttatgatgtgcctaaaagaggagagaagatgcgaagacagaa  
ttttcatatttgaaagggttcgatatcgatattgggaaacgaatcaaggtcaaaaaa  
ctcagttcaatagttgaaatttaaaaaattttattaattcaatccgattgggttcgtt  
ttgttatgggttcggttctatatcatcaaaccaatcgggttggtcctaaagataatta  
taaataattcaccaacaccagtggttaaacacatatcaacaaacctaaagttagataaa  
caaagagacccggg

**Arabidopsis Transmembrane Protein from Arabidopsis  
(clone#6468048)**

cccgggaattggcactcttcttctgctgggttccaaaagaaacgaatcaatatgtgc  
aacaagaagagctccagaagcagtcattctctaaaatcttaatacaacaacagctca  
agaagaaaaaaattccatagctagagagaaacaaaagtcaagaagcagcgtcgtaga  
ggcacaagaatcaaacctgaatggcttaagccgaactgagtggttttgactagaccat  
catcagaaaagtctccaagacggtagtcggatggttagatcgctcaagtaatttttgg  
ttttgttggtctcacgttttcagctgcccatttgatttcagtttgggcttttcccta  
tctctaaaggcccaatttcatttaggtttagtttatttgatcattatccttactata  
aaggcttcgccttccgagaaatttaggggttcttctgtctgtctcgtcactcaggtt  
tgtcctcaacgactgcttcaactctagcttgattcttcttcttctcgttttatgtat  
actgtacattagattattcttgtttctcgagcttctgctatagattttgattctttt  
ttttgttgctttgtttcgtttccaggatcagatcttagctaaattgagacaagctc  
aaaatgaggtacttgacgcatctcttacattcactgtttaattagagaacaatacgt  
ctctgaatcgtgattcagagacgtattgttcttctgtcatatgcaataagtttaatt  
agagaacaatacgtctctgaatcgtgattgttttttggatgtgcgttattgatagct

61

ttatgatgttaatagtctaggattgacacgaagttgttctgcagttttgcataaaatg  
ctctttactaaggcctctaaatttggatgacaaatctaaatcttgccctataaaaaat  
ttaggtgtattaagataagattatctttgtatggtagtgtctataatgtgggttggtc  
atggtgaggttgtcaatgttgtgtatctttgtttgtttagtttaatttgcttaactct  
gttctttgtgggttaatacagtaagcttcagagtgaggccgttcgtgaagccatcac  
tactatcacagggaaatccgaggcaaagaaacgtaactttgtcgagactattgagct  
ccagatcgggtctgaagaactatgacctcaaaaggacaagcgtttcagtggtatctgt  
caagttaccacatatccccgcctctaaatgaagatctgcatgctcggagatgccc  
gcatgttgaagaggtgatatactctttcatggaaattgatcattttgtgctctgttt  
cttgtataatgggtttgtgctcatttcatttgggtggtctattagtttcatttgatg  
ttgtatatgtcttctgaatgttagatgcatgatgttttcggaatttggtcattgttta  
tttaggcttcatttctgcataattaaatatttgcttatttcattctgtatcttttc  
gtaggctgagaagatggggttggaaaacatggatgttgagctctctaaaaagcttaa  
caagaacaagaaactcgtcaagaagcttgcaagaaataccatgctttcttgccctc  
tgagttctgtcatttaagcagattcctcgtcttcttggtcctggtcttaacaaggcagg  
caagttctggctacagctaataattccattgttcttctttacatccgttttgatttg  
gataggttttagtagtctatttctttgtcaatgtcttttgatacaatgccaatcc  
tttatcctgtgagattatgcttctttgatgattccttaagtaacattcctttgcttta  
ctttacacaggaaaattcccaactcttgtagccaccaggaatccttgaggtcaaag  
gtgaatgaaacaaaggcaacagtgaagttccagctgaagaaggttctgtgcatggga  
gttgaggttggttaacctttcccggt

**Diaminopimelate Decarboxylase from Arabidopsis  
(clone#4159709)**

cccggtggcaaaactgagatataagaggggaaggtgattttcatgcaaatttttttt  
tatttttttttgaaatgaatgcaaaatttattcaaaaaaaaaaacctgggtacatc  
aagtacttctatttctgagtttttgaaaaatctaaagacaacaaaagactttacaatt  
taataaaaaataataaaaaatactttatcactctcaacgaaattgttgatttaataa  
cgtatctcttgggtaaaaacagcgtttttatttgacgaaattgttataaatgaataaaat  
gataatagaaactagtgtggtacgtaaaaataacctctcatttggcaaaataacggtta  
tgtatcatgagttatgcatagcagcgtgcttaaatagtgtgctttcaggagaaaa  
tatataccaagttatttgtgctgaaattaccacgcaaatctgaggttcgaatggcaaaa  
taaaaaaccaatgtcatttctttaatgtatttaaggtcatttaataaaaattgtacac  
ttttttcacctgtaagcgttccaaagtgtagaatggataactagaaggggtcaaaggt  
ataatattaataagcgaactcactttttgcccagtgatttcacttcttacatttgc  
ttgatatagttaccctaaagtgtatataattcccttatacaattgttctattttct  
ggattataaggggaataaagaaaaagaaaaagagaggtatataataataactttata  
aagtgtgttagattctaatttgaacgaaaagtccaaagtgaagaaaaaacgaaa  
aagtttttctgttttgttttatatctatagccaagaaagtctcagatttacaaga  
agtttaactgagaaaaacaaaaaaaacttatgaagcatgaaagactaattaacgag  
gtgattaattttgagacaaattaaacatcgaattaaaagtaacatttggagggttta  
tatgttatatatgtgacatgataagtcgattcatgactaatgtatatctggaatct  
aacatggaagaatagagaacgaagcagagccaaggtcaacttgccagacacgaatca  
acagattgtgaatgagaccaaatcaatgggtcataaaccggttgggtttaaacgggca  
agtcactccttgggtcaattccattcgttattccttcatgcaagacctctgatacaa  
ccaaagactcccattacaatatcttctcgatcacgagctacttattttcaaatgtgt  
tacctcttctgtagctcttgtgtgtgtggttaaagcctagtgcagatgtgtcgggtat  
atataggcatatatacaaatgcgacaaaataagtatattatattgtttaatttct  
atattccatttctatatgcatggctgggtttttgaccaaaccctaattcaagaat  
agaatccaaaagatgggatcaaagaatataatctaattgggtgaccacattttccga  
tttaattcgcatagttaatatctttccactactttatgccgcagaaatttgaatt  
aagtaagacaaagaaatacagatataagatggtcgtagaaccagtagaggaatttc  
attttctggtgataagtggaatattaataagagaatggtccttactctttacagtgg  
gaaatgggaatagtagccattataatttccatcagattctatatatgcatgtttgta  
taagctaaaaataaatacgttttaagcattctcaaaaaaatttacaagttctagagac  
tctcttaacgtcggcaatttatattctactttacatgacactttcaggaaaagaaaa  
ctatactcactagcagatcattaaattttctttttctttttttgaatgaaccttagt  
tgtggtttttatttttgttagctagaaacttcagtggtttttttccgccaatggtag

62

tgctttgatgatgggtccggcccg

**Peroxidase from Arabidopsis (clone#4006885)**

cccggaatcaaggtaacgaaggagatcagcgaaaggatgggctatatattggagt  
tttttctgctgtaagtaagtctttgtgatcttccatgaggacataaactgaaga  
ataaactcaactcattgtgttctggtgtgtttcttctgatcagattcctcggtgcat  
ctgcaacttttctgctgtgggggctttatttataaaacaagagtagagcgtgtggttaa  
tcttcataatcttctacaattccacttccattctctaattattctctcacgtgat  
acacacactcaatcactgatgtactcgtatggatgcagcgtggaactgatgcattgc  
cggggatgtcacttctatcggttactagaaactgttaagtattacaagaaaactca  
aaaggattccatttatgcaaaatctaagagaaaagctcactgtggtctttggttaca  
tttatggatctctcaagagacaaatgctatgtaagctaattgattttggtcttgata  
aacaggtgagtggaagtggacaaagctactcaagaactgaagacatcaacaatgctt  
ttgccaatgaagtctcatgggaccgctcttccgcatcttctactcaagcgacaacaa  
cacagagaccaagtgaagaacataggtgcgatctaattttgtcaagtgcctcaca  
agaggtactgtttcaagccatgggtatggcagcgttggtgatctgcgatttctggatt  
tgctttgtatgtttatttctacttctagaaagaggtcaaaaagttaatagcttca  
ccgtgagaatgttggtttcaccagattcatgtgctatgatagaaaagacaaagcaa  
acaagagtcttttctttgcttaggttacaagaacaagagtatcgttataaagtcaac  
aaagattgaacataattttgtcaagggagtggttagaatctcttctactctcttg  
cctttctcactaagacaaaaaaagacttggactttgtctaagggtttgtggatatt  
attaaccaagtcttttgcaaaaagtaataattgtttttcgcattcctcttttagaa  
tttagtttaattctaggcttataattggttattactttcttgaaaaatgatctgttta  
ttctattcatacttgggttacctcgctttttatcttacttctacaaaaggattatcag  
tgaaagttagtctcttactctcaccttccgaaaaataaaacaaaaatctcgatactt  
tagatcaaaccaagttgattaaaacatccctattccctacgattctgatcttgagat  
atattatcatgttaagatctaaattgacaagaaaactgatttttctattcttagagg  
aaaaataattactattagtcatgattgtcgaccgtaagaggtggttttagttact  
ctccatctttctttgaagaagtcaagaagtcaagaattatatcaaattaaacatcaa  
tattgaacacatatatctgtatgggtttatgtttagaaaaattccaatatttatatat  
tcttagggaaaaagaagcttattcttcaaattattgttatgagtcgttaaaatagtg  
ataaaaatataaagtctaaattataaaaactcagtttgcttttgcttttaccctctca  
agtctccaaagtcaaattaatttttagttaattaaacaaaaaaaggtttattagtcaa  
acttagcatgcaatgctgggtaccaaaccacagcattagtctcttttaattctcttt  
ttctccaataagtttttacaatttttaattgtttgcatttcccttgattatttatct  
tcacccaatttagctaataccaactcggtttcttattcttccaagtccttttctat  
aaatacgttcttctccctcttatttcatatcactcaccacaaagtccttctcatt  
cctcateccggg

**Mitochondrial Uncoupler from Arabidopsis**

(clone#4220510)

cccggaatggtgtgagtgaggagaagaagagggaaacaaaggatatttattttagc  
gagttttgtttgtgacgcggttttgtctgtgttcaatgttgacgaaacgagtgaga  
gagtgctgattattaaagaaaaaccctaattaaagtcagacccgcggttatataaaat  
agtcaaaaagtaggaaaacgcgtgtgtgagtgagacagagacagccattgtttgct  
ttatgggcttataagcgagacgtgttaattgggctttttcctttatggccgaaaaca  
aaagaaacgtcgctgagagattcgaactctcgcgggcagagcccatgtacttagca  
ggcacacgccttaaccactcggccaaagcgacttgttgctatgagttagacaaaatc  
attaaaattctctattatgatttctcatagtgtgtgtgtatattgtggatctactaa  
aaattctttgttattattactttattttgtgaattagtttgatataggttaagtacaa  
agtttaactttattatttactcaaaatttatcagattaactgattttatattgtttcc  
tttggtatatagacgtactatagtttttagaaaaaccataagattccttttatatttc  
atagagtgaagagatgagatgagatcttggctggagaagaaataagtttccacgagg  
aggactcttttttttgggtgaagcaggaggagactcttggttgatccagtcctt  
acgttagacatcgacccctacattttatttgctttctctatcaacatggcaggtaaa  
aatcttctattcaaccgaaccaaccaaagtccttcccaataatattcaagcaccatc  
ctttgggaaactcatacatactacagtctacactctttcattttctttcaacgctca  
acttaacaaatgatatagtttagttgtcaattatatgttttaattagttgtttcaca

63

tcaaattctggtttgatatttgatgactattttcggaacatctcaatgtcccgcaa  
atacaatcgtctatcatatataatcccgtagcttgattcttatagatagaataata  
tggcgtgatctttataatataacatataagaatcgtgtagatttattttattttatt  
ttatatatcgcataaattgcaaaatacttatatatgtttgttatatatgatacccat  
tttatagttacttaaaaaaagttaagcgataatatatatatcaactttttataac  
aaaaaggtataacacatgggtaaagaaaaataaaaaatgaagacatgggtgtgacacgaa  
aatggcactaaatatacatatataatagatagctacaatatcccatcatacacactt  
tttaattgactaatacacaacttacacacttttttaattgactaattcataacttt  
ttatcattgtcaacatgcaaattcatatttccggtgaactattattcttattttgtt  
tttaaaagaagggtcttctggtaataaaaaatagatttccaaatgacgttagagcaa  
aaaaaaaaaaaaagggtgtctgggtctggtaaaatgaaaaagcaaagcgtcttggtatag  
aaaagtaataatactgcctcctaatttcttctgctcttaccgaagaatctctccact  
cttgcctctcttgcgaaccctaaaccagaagcaccagattttttcaactttttccca  
gagaacaatagaaaacccaacttgtgctcttaggggttttcttattcttctcctc  
tttggtatttcttgggtcatcattttggaagcttaccaccagcgaaaaaattataa  
cttccatcgattcctggcttctctctcgtctctctgcatgtgctaaatcgccgg  
actgatcctcactgtcacctctgttcccggg

**Stress protein from Arabidopsis (clone#6598614)**

ccccgggattaggggtttgagttgtcactggaaagagggtttgattgtgagtgatgat  
ggagagattatgaaggagtttgtgtgtatttatagaggagttagggttttgagggtt  
gatgagaagtaggtttgaagaagttttgttgttgcaacttatttagagttacttgtt  
ccacaaccacaagtaagattgggtcacttctaagttctaactagaaacaacctgaca  
catggagatttcagctaacctagtttaattgtatatgtattatattttatttaaata  
tataaaataaaaataaattttcacaataaaaagaactacaaaaaagtgagaaaaataa  
tttgataaacaatttagaaaaattagtatatcaataaaataaatttataatccgatgg  
ttttgccttttgggtttggcctttgtttgaacttcgatgagtgactatgtatagcgaa  
aacaattcgggtttgtttttgggtttaattttaaaaaatacaagcgacaatatctgatg  
agaatagggtgaaaagcaaataatatcagtttaattggaaatatctacttttttaca  
ctaatttttgttttgggtcaaccaacaatatagatttaatttaattatgggtttatgagct  
tttatttgggtgcgacagtatatatatgtttaaatagtgatattgcatggcggaagggt  
cgggaagcaacacatatctcctttttaaattttttttaacaagaataaacatgttaa  
tttttttttgaaattaataaagaatacatatttctaatttttgcgtcagatagatga  
ttaaagagtggtgtgttttttttaacaacaaggaatacattatacatatttcatatt  
tctctcgacattgtttgttttttaaaaaatagattaaagagttctacgaagctaagt  
agctaacgaagacttgaaatgagaagaagacgagaatcttttaatatttttgttaa  
gcgataatattttgaaatttaataaatatagatttaagggaataaacaataacgcagat  
atcggtaagtcatagaaaaaaagaaacaacacaaacttacataaacatgtttcctaa  
tttgtaattggagtaaaattccttcttttttttttttttttgatttggattccaatta  
gtaagaactcaatgactataaataacctttaaccctctcattatttcttactatca  
attgattaagctctcgttcctaagaaagcaatagacgaacaagaacccatcgaagaa  
cacaatatctcttttgaagttgtcgataatgttagtacaccgttacttcgtccaaga  
cttttttgcggttcggtttcttacaacaaggatttgggttaccattacttttgtcg  
taactcctttttacatgtacgtcaaaaagtggttctcgtcctcggttgaagaaacg  
accttcttaccacaaaaagcttattttaaaccttcaaaaccggaaaatctcaatc  
taaaccggatcaggttcagtgagaaaccgattcaaacaccgagtgagaagtagaatt  
tttgatgggttcggtcacatgtgtgctgctccttcgccaagacatgtaccgattcc  
gatatttttgggtgtaaagatgatcaaagagtttcaaagctaagcagcacttgaat  
gagaagaagaagaccaattactcaattagattttgttttgggagcaattattgtct  
atttatctttgttttagcaataatctgtatccactaatcttcacagtacttgact  
aacaagaagtaaaagagttttcttatttccaattgttttttaactctgatacttttttc  
ataattttacaatgtttgatgaaaaaaacattcaaacctaaattttctttttttgg  
tatgaattcaaacctgaattacttttgacgaggacccgacggtataaatagggtgat  
ctccaacaacaacaaaagggtcccggg

**Pectinacetylsterase from Arabidopsis**

(clone#6671954)

ccccgggtggtggggacaatggatccgggtctgcgtagcaacaaggctgaaaaagatta




FIG. 1




FIG. 2




FIG. 3




FIG. 4



# Selectable Markers

pNOS / NPT-II / tNOS

pSU / Bar / tNOS

pSU/ Intron / Bar / tNOS

pUBQ3 / Intron / PMI / tNOS





FIG. 5

## Scorable Markers



<sup>1</sup> Construct useful for promoter analysis.

<sup>2</sup> Construct useful for high constitutive expression of genes of interest.

FIG. 6




FIG. 7




FIG. 8

AKK110P1  
SEQUENCE LISTING

<110> Mushegian, Arcady R.  
Taylor, Christopher G.  
Feitelson, Gerald S.  
Eroshkin, Alexey M.

<120> Materials and Methods for RNAi Control of Nematodes

<130> AKK-110P

<140>

<141>

<160> 139

<170> PatentIn Ver. 2.1

<210> 1

<211> 165

<212> DNA

<213> *Globodera rostochiensis*

<400> 1

```
gtttgagatt attgactttg catattttcca accaagttca tttgaccaat attttcctgc 60
taaaccatagc aaaaatggtg aaaccgaagg tcggcattaa tggctttgga cgcattgggc 120
gcttggcgtt gcgcgctgcg gttgagaagg acaccgttca ggtgg 165
```

<210> 2

<211> 342

<212> DNA

<213> *Globodera rostochiensis*

<400> 2

```
cgactacatg gtatacatgt tcaactacga ctcgacccat ggccgcttca atggcaaaaat 60
ttcgacaagc gccggcaatt tggtcgttga gaaagagggg aaggccacgc acaccatcaa 120
ggtgttcaac ctcaaggacc cggccgagat caaatgggct gaggtgggcg cggaatatgt 180
gatcgagtc accgggggtgt tcaactacat tgagaaggct tcggcacact tgaagggggg 240
cgccaagaag gtggtcatct ctgctccgct cgctgatgca ccgatgtacg tgatgggcgt 300
caacgaggac aaatatgacc cggccaagga caacgtgatt ag 342
```

<210> 3

<211> 205

<212> DNA

<213> *Globodera rostochiensis*

<400> 3

```
gaagccggcc tcattggacg ccatcaaggc ggcggtgaag aaggctgccg aagggaattt 60
gaagggcatt ttgggttaca cagaggacca ggtggtgtcc acggactttc ttggagacag 120
tcgctcgtcg atcttcgacg ctggggcggtg catctcgttg aaccgcact ttgtcaagtt 180
ggtcagctgg tacgacaatg aattt 205
```

<210> 4

<211> 167

<212> DNA

<213> *Globodera rostochiensis*

<400> 4

```
ttaaaccgatt tattcacacg cacggagaaa tgaggattac ctaatttgat tgagtccttc 60
tcgtccattt gtcaattgtg gccctaaaga gggccggttg ggtagtttt ttggtgttcc 120
ttctccttgc tggctcaacc accgaagccg tacagcgtcc ggccttg 167
```

<210> 5

## AKK110P1

<211> 41  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 5  
 catggccgtc acggtcttgc gcttcgcgtg ttcgcagtat g 41

<210> 6  
 <211> 79  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 6  
 gtttcccagg aaaactttca gcacggaacg agtctcctcg taaatgaggc cagagatgcg 60  
 cttaacgcct ccacgacgg 79

<210> 7  
 <211> 168  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 7  
 cggcttggtg atgccctgga tgttatcccg caagactttt cggtggcgtt ttgcgcctcc 60  
 ctttccgagt ctttttccgc cttttccgcg tccggacatt ttgttggtta atcagaagag 120  
 cacagagagt aggagaaata ggaaattttg cctcgtgccg aacgtgcc 168

<210> 8  
 <211> 330  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 8  
 gacagtctcc gttctggtta tgtgtcacac gcgcgaactt gctttccaaa tttctaagga 60  
 atacgagcga ttcaccaagt acatgccggg agtgaagggt tccgtattct tcggagggat 120  
 gccgataaag aaagacgaag aggtattggc taagaacacg ccgcacattg tcgtcggaac 180  
 gccgggacgt cttttggcct taggacgcac tggacatctg aagctgaaag gcgtcaaatc 240  
 ctttgtgctg gacgaatgcg acaaaatgat tggagatgcc gacatgcgcc acgacgtgca 300  
 ggaaatcttc aaaatgacgc ctcaggagaa 330

<210> 9  
 <211> 136  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 9  
 actttgccgc gggagctgcg cgtcttctgc aaaaagttca tgcaggaacc aatggaggtg 60  
 tacgtcgacg acgaggctaa gcttacgctt cacggctctc aacaatacta cgtagactg 120  
 aaggaaaatg agaaga 136

<210> 10  
 <211> 141  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 10  
 tattaaaata aaatacaaac aataatataa tggctgtttt ttctgtcatg tttcaagttt 60  
 ttgtgttca tcactttctt cagcagcgac aatacggcca atccggtgaa agggccaag 120  
 tcaatagctc gctcgtacc t 141

<210> 11  
 <211> 141  
 <212> DNA

AKK110P1

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 11

acccaggcac tctgttcac ttcggcatcg ctttttggca atgtcaacaa cactttgctg 60  
 gccattttgt ttctacagca cagcacacc gtcgtcttta cagcgttcac ctgccaataa 120  
 aagtagccgt atttgcgaaa t 141

&lt;210&gt; 12

&lt;211&gt; 37

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 12

gcgttgggtg caagctgtac acaaggtcgc ccggttt 37

&lt;210&gt; 13

&lt;211&gt; 161

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 13

gcgcgttcca tcgcccgcac cacaaaaagt cccatcgctt catatcgtag cgcaaattgt 60  
 ctttgggtgca aatggcaaaa cggccaaaat aatgggtcgaa gccgtacaca accgccaccg 120  
 ccacagcgcc aaccacacac caaatgcgaa atttatcgaa a 161

&lt;210&gt; 14

&lt;211&gt; 306

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 14

gaattcgttt gaggtataaa taaataataa atggcagcca acgaatcgct aaatgtggac 60  
 agtttgatca ctcgattgtt agaagttcgg gggtgtagac cgggaaaaac agtgcaaatg 120  
 gacgaatctg agatacgcac ttgtgcatc aaaacacgtg aaattttgct gtcgcagcca 180  
 atcttgttgg agctcgaggc acctttaaaa atttgtggtg acattcacgg acaatataat 240  
 gatcttctga gattgttcga atatggtggg tttccaccgg aagcgaacta tctatttctt 300  
 ggggac 306

&lt;210&gt; 15

&lt;211&gt; 261

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 15

gcaaagcctt gagacgattt gtttgcgtct tgcttacaag attaaatatt ctgaaaattt 60  
 ttttcttctt cgtggcaatc acgaatgtgc ttcaatcaat cggatttacg gattttatga 120  
 tgaatgcaaa cggagggttc tcaatcaagt tgtggaagac cttactgac tgcttcaact 180  
 gtctgccaat tgccgcttta atcgacgaaa agatcttttg ctgccacgga ggctgtctcc 240  
 tgatttgcta aacatggcag c 261

&lt;210&gt; 16

&lt;211&gt; 151

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 16

gaattctttg agtgcattca gcgtttaatt ttttcgtatt ataataagca tggctcgagg 60  
 acccaaaaag catttgaagc gacttgcagc acccaaaaaa tggatgttgg acaaattggg 120  
 tggcgttttt gcgccacgtc cattgtgcgg a 151

&lt;210&gt; 17

&lt;211&gt; 306

AKK110P1

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 17

```

tcaagtacgc gcagtcgtac aacgaagcgc gcatgatctg caaacagcgg ctgatcaagg 60
tggacggcaa agtgcgcacc gagatgcgct tcccgtgcgg aataatggat gtgatctcga 120
ttgagaagac aaacgaaacg ttctgtctgg tgtacgatgt gaagggccgt ttgtcatcc 180
atcgaattca aaagctggag ggccagtaca agctgtgcaa agtgaagaag caggccgtcg 240
gggacaagca ggtcccctac attgtcacac atgacgcgcg caccattcgc taccggaccg 300
ctcatc                                     306

```

&lt;210&gt; 18

&lt;211&gt; 528

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 18

```

gaattcgcac aacgaattga agacttatgc ggcagaaaaa ggacttttgc caaagtgtga 60
ggagcaagca gacgaccttt cggattggct ttgttcgtcc attgggttgg agcatcgccc 120
gttcctaccg tatacaaacg ctgtaataaa tgaaacaatt cgattagtca atttgatccc 180
gttcaatctt agccatttgg cgcttgaaga tatgcaaat ggcaatttta ttgtgaagcg 240
tgggacacca attgtaccgc aggtcagcag tgttctgttc gacgaaaaac tgtatccgga 300
gcccgatcgg tttttgcccg aacgctttct ggacgatgag ggccgtttga agaaaagcga 360
cgaacttatt gcatttgggg ttgggaaaag gcaatgtgcc ggcgaaagctt tggcccgaat 420
gacacttttt ctgtttgccc ctaatttctt tctcgcctac aaagtctctc cgtccgatcc 480
actgaatcct ccaagcctga aaaagttggc ggattatctg ttacaca          528

```

&lt;210&gt; 19

&lt;211&gt; 335

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 19

```

gaattctttg agaaagcggg aattcgtttt tggctataaa atgattctgt gggccacgat 60
tttgttgatg gctttggaca ttgcgttcgg tggcaccaat caaatggaat ttgatcagtc 120
ggcgcgcatg ttccccgact cccagttcat cgatttgatt tcgcgcgaca tcgaatcctt 180
ctccgcccga ttgggcgttg gccataaatt tatgagcggc ggtgccggtg agggcgcca 240
acagctaggc cccgaggggc cctttgagca gcggcaacag gtgaagagtg acaatgttct 300
ccccgcgtat tgcgagcctc caaatccctg tccga          335

```

&lt;210&gt; 20

&lt;211&gt; 52

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 20

```

ggacggctgc acggaacagt tcgagaacac tgccgagttt tcgcgcagct ac          52

```

&lt;210&gt; 21

&lt;211&gt; 190

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 21

```

gcttgtgtga ccaggagcac atgtttaact gtccgtcgaa gaacaaccgc gaggagtacg 60
agcaggatct ggagcaattg ctggccaaca acggactgca caaatcaatg attgccaaga 120
aattccatct cacgcgggcg gaggagccgc gccgtcgaaa acgctcttgt cgcccggctt 180
cggccaaccg                                     190

```

&lt;210&gt; 22

&lt;211&gt; 52

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis



## AKK110P1

<400> 22  
ccgtacaac ccctacctgg agggcgcccc gctgaagtca gtggccaaaa ag 52

<210> 23  
<211> 54  
<212> DNA  
<213> Globodera rostochiensis

<400> 23  
gaattccgac tctcaaggtg gacccacgcc caaccaacag caattgtcag ctgc 54

<210> 24  
<211> 77  
<212> DNA  
<213> Globodera rostochiensis

<400> 24  
ccgcacatgt cgaggcctcc atcttttggc actggtcatc accttccgcc tactgctaac 60  
aacagaccgg aacagca 77

<210> 25  
<211> 439  
<212> DNA  
<213> Globodera rostochiensis

<400> 25  
gtcaatcaaa aacgccgact tcgattcctc agctgatggt cagtaatgcg ctaaagggca 60  
tccattccgt ctctttctaca tcagcaacac aatcacattc cacgcccagt tttatgacac 120  
acaacgtgca gcagcaacat gttgttggtc aacaacagca gcaacaacag aatttccaac 180  
aaccgccgcc cctatcgtae actcacagcc accaacaaca aaaacaacca ccacaagcgt 240  
cacagtcgat gttgtcaatg aaaagtggca atgtttgtcgt tgtttgttccg caacaatcgc 300  
agcagcacca ctaccaacag cggacactga cgccactgaa gcacacatcc gcctcctcca 360  
cgtccgatcg cttcgtcatc accaaaacca acagggtgct tccactcccg tcgcagcaag 420  
gcgccacggc cactgatga 439

<210> 26  
<211> 539  
<212> DNA  
<213> Globodera rostochiensis

<400> 26  
gaattcgttt gagacacatc caattaatta atagttattg ttggcaatgg gacgagttcg 60  
aacgaaaact gtgaagaagg cgtcgcgcgt cattattgag aagtattaca ccaaattggg 120  
cctcgacttt cacaccaaca agcgcatttg cgaggaggtg gccattatcc caagcaaacg 180  
gatgcggaac cgaattgcgg gattttatcac acatctgatg aagcgcattg agctggggccc 240  
tgtccgtggc atttccatca aattgcagga ggaggagcgc gagcgtcgcg acaattacat 300  
gcccgaatc tcttacctgg atgcgcagaa tcaccagatg atcagcaccg accaagagac 360  
gaaggatatg gcggaatttc tggggctagg cctcaacttg gaagtgaag ggcctttgac 420  
gagtggcggc gctggcgag gacgtcgttg agtcaggaca attggcatta ttgttgaaaa 480  
atcatcgatg ttttgttcgc atttgatga taatgcgctg ataaattttt gttgatttt 539

<210> 27  
<211> 179  
<212> DNA  
<213> Globodera rostochiensis

<400> 27  
gaattcnaca gtttctgtga gtaatggcat ntcacactgc cggcatccaa cagtgtcttg 60  
cggccgaaaa gcgtgcggca gaaaagatta atgatgcccc gaagcgaaaa gcacagcgac 120  
ttaagcaggc caaacaagaa gcccaggcgg agatcgagca gtatcgnacg gagagggag 179

## AKK110P1

<210> 28  
 <211> 133  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 28  
 gcaaaattat ttgggcacgc gcgacgacat cgagcagcaa ataaagcgcg agacagaaga 60  
 gtcgctggag gcaatgaatc gcaatgtcgc ggcgaacaaa cagcaggcca ttgtacgtct 120  
 gctgcagttg gtg 133

<210> 29  
 <211> 482  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 29  
 gaattcgtga aatcaaaagc ttttttaatt tatttacaca aaaaatgggt ccaccaccaa 60  
 ttgcgctgtt ggtcactggt gccgctggac aaattggcta ttcactgggt ctgcaaatcg 120  
 caaaaaggcga tgtgtttggc aaagatcagc caattgttct cgttctcttc gacattccac 180  
 cgatggccga agtactctct ggtgtccatt ttgaattgat ggactgtgcy ttggcaaac 240  
 ttgccggtgt ggaggctgtg accacggaag agcaggcctt caaggacatt gactacgctt 300  
 ttctgtcgg agcgatgccc cgaagagagg gaatggaacg aaaggacctt ttggcggcaa 360  
 atgtcaaaat tttcaagtcc caaggcgaag cattggcccg cttttccaag cccgtncgtc 420  
 aaagtctctg tgggtgggcaa cccggccaac acgaacgcgt acatttgcgc aaaatatgcc 480  
 gg 482

<210> 30  
 <211> 605  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 30  
 gaattcaaaag tgccgaaagc gttcaaaata attccggcaa tggcaaatg ggaacaaatt 60  
 ctagacctta cttccccga aaaatggagt tcagcggcga tgtttcaagc aactcgtgtg 120  
 ttttctgcca ccggcacacc gtcacaatgc caaagggtta acactttggt gctgttgcca 180  
 cgactccgtg atgagattga cgagtacaag aagctaaact ttcatttgta tcagtgttg 240  
 tttaaagcaa tgttcaagcc ggccggattt tttaaaggca ttattttgcc tctttgcaa 300  
 tctggcactt gactctccg tgaagccatc atctttgggt ctgctctgcy aaagatttca 360  
 ataccgaac tccacgcccgc tgcagcaatg ctcagcatag caaaaatgga ctactcgggc 420  
 gccatttctt ttatcctacg tgttcttggt gaaaaaaatt acacacttcc tttccgagca 480  
 ttagacggcc tcgtttttca ttttcttgga atgcgctcac atcagggcga gctgccagt 540  
 atttggcacc agacactgtt ggcttttgct gagcgttacg caaaagacat aagtgcagaa 600  
 cagag 605

<210> 31  
 <211> 112  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 31  
 ccattcccat catcaaatta ccccgattta ctgcggcttt tgccggcgcc cgagtcgagg 60  
 aatgaggaaa gtgaagcaaa tgtgcccgtt tatgcgcgta atgatgaaat gg 112

<210> 32  
 <211> 105  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 32  
 gaattcgttt gagcatttat ttgacaaaat ctgaataaat ggccgtacca aaagaagtta 60  
 ttgacaaaat cgaggcggtt tacaagaagc ttcaggaagc gtctn 105

<210> 33

AKK110P1

&lt;211&gt; 425

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 33

```

aagaagtacc tcaccaagga agtcgtcgat gcctgcaagg ataagcgcac caagcttggg 60
gcgaccttgc tggatgtgat ccagtcgggc gttgccaact tggacagcgg agttgggggtg 120
tacgctcctg acgctgaggg ttacaccttg ttcaagccgt tggtcgaccc gatcatcaac 180
gactaccatg gtggctttgg tccgggcagc aagcagccgg caactgacct tggtagcggc 240
aaaacgcana tgctgaccgg atctcgaccc cgaggggaaa atttatcaat ttcgacacgc 300
gttcgtttgcg gccgtttcct ttaagggata cccggttcaa cccgtgcttg acnaaaggan 360
aactacnttt ggagatggga aacnaaggtc nagggccgtt ttctaacatt ttnaagggcn 420
atcct 425

```

&lt;210&gt; 34

&lt;211&gt; 581

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 34

```

gaattcgttt gagcgaagag ttttgtggtt gacaccgggt tatggacttt tagcccgtga 60
tccttgacgg tccaaagccg cgttcagttc cgtgccgtgt tttttaaag aggcgggagag 120
tttgacggtc attccaagca gccataaac caccaaaacc aaataccccc cccaatcga 180
tccccccct ccaattctc cgcattatc gcattatcaa ttctaatacag cacaaccact 240
gcatcattcc tttcccgacc atacgatgct aagtgaact ttgaaaattg gttcatcgg 300
agccggaaaag atggcccaag cattggcaag aggacttatc aattcggggc gataccggc 360
agagaatttg atggcgagtt gtccaaagac ggacgaggct ttactggagc aatgcaaaaa 420
attgggaatc ggaacgacgc acgacaacac ttgggtcgcg cgagagaacg acgtcatcgt 480
attggcggtc aagccgatgc acatcagcaa agtgacgtcg gaaatcgcac ccaatttcg 540
gaggggaacat ttgcttattt cattgattag gaattacact t 581

```

&lt;210&gt; 35

&lt;211&gt; 102

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 35

```

gaattcgttt gagaatttta ctttatataa ttgacgttta atcagcagcc ataagcaatg 60
cccatcaaa gcatccggaga aacattaagg aagtttattg tc 102

```

&lt;210&gt; 36

&lt;211&gt; 34

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 36

```

tgcaaatgat gcaaacccca cgcttcacaa gatg 34

```

&lt;210&gt; 37

&lt;211&gt; 100

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 37

```

tcattgtgtg gccaaatctc gcttctggta ctttacgagc atgctgcgtc gagttaagaa 60
aacacacgga gagatcgttt cgtgtcaaga ggttttcgag 100

```

&lt;210&gt; 38

&lt;211&gt; 176

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 38

## AKK110P1

tgaagaactt cggaatttgg ctccgttacg attctcgtac tggacaccac aatatgtacc 60  
 gcgagtatcg ctgatgttac cgaggccggt gccgtgacct aatgctatcg cgacatgggc 120  
 gctcgtcacc gcgctcaggc ggatcgaatt caaatcatca aagtgcacaa ctcaag 176

<210> 39  
 <211> 155  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 39  
 gaattccaag tttgaggtat tgtttgttat acgatttctt acaaatgaca gaacaaactg 60  
 agcgcgcgtt ccaaaaacaa ccgatcggtt ttctgaacga caagttcaga acgcaaggga 120  
 ttgggaagaa ggcattcaac aaggaccgtt actgg 155

<210> 40  
 <211> 35  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 40  
 tcctcgcgag gctattgagg gcatatatat cgaca 35

<210> 41  
 <211> 70  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 41  
 tggaaatgtg tccatccgag gtcgcattct cactgggggtg gtgatcaaaa acaaaatgca 60  
 gcggacgatt 70

<210> 42  
 <211> 85  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 42  
 tcgtacaaa atatcgctgc tatgagaaac gccacaaaaa catgtccgtc cactgttcgc 60  
 cgtgttccg agatgtctct ctgg 85

<210> 43  
 <211> 193  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 43  
 agttcggttc aatgtgctca aggtgatcaa agcatcgggc tcgaagaaag cgttcgacaa 60  
 attctgagtc ggccaagcca accgcgaacg gtcatttggt atggttccta attgttgctg 120  
 tttttcaatt atttgtgta aatgactgaa ttatgatca acggtatact agtattcttc 180  
 tgaaaaagct cga 193

<210> 44  
 <211> 219  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 44  
 gaattcattt agatttggtt tgaagctaga aatctttatt ttgggagtc acaacaatgg 60  
 gaagacgtcc ggcgcggtgt tatcgctata ttaagaacaa gccgtatccg aagtcgcgct 120  
 ttgtcgcgg tgcacccgac ccaaaaattc gcatttttga ttggggtaga aagcgcgcca 180  
 ccggtgacga attcccatgc tgcgtgcata tgatcga 219

## AKK110P1

<210> 45  
 <211> 489  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 45  
 tccgaggcgc ttgaggctgc gcaatttgt gcaacaaat atatggtgaa gaattgcgga 60  
 aaggacgggt ttcatatgcy cgtcagaatc catccatacc atgtaattcg catcaacaaa 120  
 atgttgtcct ggcgtggtgc ggaccgtctg cagactggga tgcgtggtgc gttcggaaaag 180  
 cctcaggggac tcgtggcgcg tgtcagcatc ggtgatatgc tgatgtcagt gcgtattcgt 240  
 gaccaacacc aagctcacgc attggaggcg ttccgtcggg ctaaattcaa gttccctggt 300  
 cgtcaataca tcgtcttgtc ccgcaagtgg ggcttcacca aattcgatcg cgaggatatac 360  
 gagaaatacc gcaaggaggg ccgtgttatc cctgacggtg tgcattgcaa gttactcaag 420  
 caacacggac ccgtgaagg agtggctcaa gaacccatt taatcttctg tttgtcttgt 480  
 gactcttgg 489

<210> 46  
 <211> 101  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 46  
 gaattccccg gctcgagccg ggttgacgat gtcctcctcc acctcctctc actgcgttcc 60  
 gtcctccttc agccggaat tgttctgtg gctgttgccg g 101

<210> 47  
 <211> 485  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 47  
 tccaccaaag tccattcgct gtcgccagtc catttattcc acaaaaagat gattccgctcg 60  
 tcgttccgat gacgtcgttt ggccaaccgt tgccccgctc accgctttca ctggtgcca 120  
 acccgccgct ttattttgtg tttccagaaa acttgccgtt ggagcggccc ttcgacgagc 180  
 aaaacgacgg ctccgaggag gaattagccg aagaagcgat gggaacgaag gcgaagaggg 240  
 cgcaaacggt cgtccgattc ggcaaaaggg cgcaaacatt tgtgcggttc ggaaagcgtg 300  
 cacaacatt tgtacgcctc ggaagggaca cgcaaaaggca attcgatggg aaaatgcaaa 360  
 gtgaacagca acagaaaaag gcttaaagca aacggcgggc acttttcttt taatgaatgc 420  
 gcgcccaccg catgacaatt cttttgtgta atgtgttgcg atttttatga tcggtaaatg 480  
 taaca 485

<210> 48  
 <211> 651  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 48  
 atctgttcaa gggactgttc ggcaagaagg aaatgcgcat tctgatggtt gggttggacg 60  
 ctgctggaaa gacgaccatt ctgtacaagt taaagctcgg cgaaattgtc accaccatcc 120  
 caacaattgg cttcaacgtg gaaaccgtcg aatacagaaa catctcgttc actgtttggg 180  
 acgtgggtgg tcaagacaaa attcgtccac tttggaggca ctacttccag aacacgcaag 240  
 gactgatctt cgtcgtggac agcaacgatc gcgagcgtgt gggcgaggcg cgtgaagagt 300  
 tgatgcgaat gctggcggag gacgagttgc gcgacgcggt gttgctgggt ttcgctaaca 360  
 aacaggattt gccgaatgcy atgaacgccg ccgaactgac agacagactt ggactgcaca 420  
 acttgcgaaa ccgcaattgg tacatccagg ccacctgcgc gacttcgggc gacggactct 480  
 acgagggact ggactggctg agcaaccagc tcaagaacag aggctaagct gggttggtgt 540  
 ctgttgcaat tgcgcgcgga attgatgacg attgaattta tttgtgtgtt tgcgcgcgca 600  
 gctcttttgt gggacgtccg attaatattg ataattattt tattccgtgt t 651

<210> 49  
 <211> 660  
 <212> DNA  
 <213> Globodera rostochiensis

## AKK110P1

<400> 49  
 gaattcccaa gtttgagatc aattcagttt cacttagaca aaaatgccgc cgaatttcga 60  
 cccaactgag atcaaaatcg tgtacctgcg ttgctgctgggt ggtgaaattg gtgcaacatc 120  
 tgcacttgca ccaaaagtgg gccacttgg attgtcgccc aaaaaaattg gtgaagacat 180  
 tgcgaaggcc acacaggact ggaaagggt taagggttacc tgcaagctga caattcagaa 240  
 tcgtgtcgcc aagatcgacg ttgtcccatc ggccgcctct ctgatcatca aagagttgag 300  
 cgaacctccg cgagaccgca aaaaagtcaa aaacgtgaag cacaatggca acctgaccat 360  
 cgagcaagtg atcaacattg cgcgtcagat gcgcccctcg tcaatcgac ggaagttgca 420  
 gggcaccgtg aaggaaaattt tgggaaccgc ccagtcgggt ggctgcacca tcgatggaca 480  
 acatccgcac gacattgtgg acgcgatcag agggggagac atcgaaatac ccgaggaata 540  
 aagaaaggac ggccgctccg atttttgtgg gacggacatt gggaatttga ggtgaatgag 600  
 ttgccaatct cattcattca tcaattgttg ttattgntgg tacggataaa tttgtaattg 660

<210> 50  
 <211> 625  
 <212> DNA  
 <213> *Globodera rostochiensis*

<400> 50  
 gtgccggaac agacgctcga ggaggttagc cgtctgcagc ggacgagctc cttgttggac 60  
 gtggcaatcc gggacggcgt cccctacccc ccactgcctc ctacaaaccg atcccccgaa 120  
 tacatgaaca tgctgacccg ctcttctcgc gtgcccatt tccgcatcta ctccggcgcc 180  
 atcggaccgt acagaccttc gttgcccgtg tacacttaca acacttaccg cggtgacttc 240  
 ccctaccgca actaccgcgg ctacaccttg gcgaatgctt actggtacga ccgatactat 300  
 tacttctcgc cgtgtgacaa acgaagcatg ttccccaccc gcttcaaaca ttgtgactat 360  
 aaagcgaacc cgcactattg gcactaccgc cacacctttt gggactatcc ctaccagggc 420  
 aaatggttcg actacgacaa cctcccaat taccggccct actacaacca tcgccttaac 480  
 ggatatgctc ggccgtatca ctaccggtcc catgctgctg cccaccggtt caattaccgc 540  
 gaagggaatg tcaggaaacg ggtctgacaa atcgaaactg tccaaattga cgtgtccgc 600  
 attcgaaaga agacgaaaaa agctt 625

<210> 51  
 <211> 402  
 <212> DNA  
 <213> *Globodera rostochiensis*

<400> 51  
 gaattccaag tttgagcaac attttgaata tgaccgaagc caaaaaactt cccgaggtgc 60  
 cggaactttt gctcaagcga cgcaaaatca gagctgcgca aaaggccgca aaagcaaga 120  
 acaaatgag ttctatcaa aaagcacgga ccaagaagggt ggaaatcttc aaaagagccg 180  
 agcagtattt ggtgagtag cgtcagaagc aacgccaatt gcttgcgctg aaactggaat 240  
 cgaagaaagt cggcaattat tatgtgccag aagagcccaa actcgctttt gtggtccgaa 300  
 tcaaaaggat caataagatt catccgcgtc ctgcgaagggt tctgcagctt ctccgcttgc 360  
 gtcagatcaa caacggcgtt ttcgtaaagt tgaacaaggc ga 402

<210> 52  
 <211> 433  
 <212> DNA  
 <213> *Globodera rostochiensis*

<400> 52  
 ccgacccgta catcgttgg ggtattccga gtcagaagat catccgtcag ttggtctaca 60  
 aacgcgggta cgccaaagag aaggacagc gcattccaat aacggataac aacattgttg 120  
 agcgcagttt gggcaagcat gacgtgattt gtgtggagga tatgatccat cagatttgga 180  
 ccggtcggac cgcacttcaa acaggtgacc aacttcctat ggcctttcaa gctgagcaac 240  
 ccggtgggag ggttcaagaa gaagtccaat cacttttgtg gagggaggcg attatggaaa 300  
 ccgagaggac caaatcaaca aattattgga aagaatgggt taatgggaag gaagcggana 360  
 aagaaaggaa attgnggcgt ttttctgttg ttgttttgac gataaattgt taactccaaa 420  
 aaaaaaaaaa aaa 433

<210> 53  
 <211> 768  
 <212> DNA

AKK110P1

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 53

```

gaattcggtt gaggtcaaac ttatttagcg tattaacaa tgtccgaagg aggagcgaaa 60
aagagtagca gcggtgccaa ggggggggtt gatgtcaaga aatttgcgat cgatcttgcg 120
tccggtggta ctgcggcggc tgtctccaaa actgttggtg ctccattga acgtgtcaaa 180
ctcttggtgc aggtgcaaga tgcctccgct cacatcactg ccgacaaacg ctacaaagcg 240
attattgacg tgcttggtccg tgtgccgaaa gagcagggtt ttctgtcact gtggcggtgg 300
aacttgcca acgttatccg ttatttcccg actcaagcgc tgaacttcgc cttcaaagac 360
acctacaaac gcatctttac ggagggactg gacaaaaaca agcagttctg gtcgttcttc 420
gtcatgaatt tggcctctgg aggtgcgggc ggcgccacgt cgctgacctt tgtttatccg 480
ctgggacttt gcccgtagcg gtttgcccg tcgatgtccg aaaagctggg tcccgcgagt 540
tcaacggttt ggcccaactgc atcgcaaaaa tcttcaagtc ggacgggtccc atcggtcttt 600
accgcggctt ctctgtctcc gtccagggca tcatcattta ccgcgccgcc tactttggat 660
gctttgacac cgcaagatg attttcgcgc cggatggcaa gcagatgaat ttcttcctca 720
catgggccat cgctcaggtc gtcaccgtgt cgtccggtgt cctctcct 768

```

&lt;210&gt; 54

&lt;211&gt; 338

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 54

```

gaattccagc agattaattg gaatggctga gaacatcgaa gagattcttg ccgaaatcga 60
cggctcccaa attgaggagt atcaacgctt tttcgacatg ttcgaccgcg gaaagaatgg 120
ttacattatg gccacccaaa ttggacaaat tatgaacgcg atggagcagg actttgacga 180
aaagaccctc cgaaaattga tccgcaagtt cgacgcggac gggtccggca aactggagtt 240
cgacgagttc tgcgcgttgg tgtacacggt ggccaacact gtggacaagg acactctgcg 300
aaaggagctg aaggaggcat tccgactctt tgacaagg 338

```

&lt;210&gt; 55

&lt;211&gt; 267

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 55

```

gaaattgcgc ccgatctcag cgacaaggat ttggaggcgg cggtcgacga aattgacgag 60
gacggcagcg ggaagatcga attcgaggag ttctgggagt tgatggcggg cgaaaccgac 120
tgagaaaaga gcaaatcgat ccaaattcaa acggaccggt cccatttcac ctccatccgt 180
ccgtcgtatt attatatatt ccagtggaat ttccccatta aaattcgggtg aaagtaaaat 240
aatttgacga aaaaaaaaaa aaaaaaa 267

```

&lt;210&gt; 56

&lt;211&gt; 597

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 56

```

gaattcgctg gacacttcgc atccggagta cagccacgag cagagcatcg accagaccag 60
catcccctac cagatgggtt cgaacaagta cgcctcgtag aaggggcatga ccggctttgg 120
acagccccgt tgggaggtgc ttgaccgctc catctcgtac cagaaccgca agtcgcaagg 180
aatgggttcgt ctacagtcgg gtaccaaccg gttcgcttcc caggcgggca tgaccggctt 240
cggcacaccc aggaacacca cctatgaggc ggaggcaggc gagctgccct acgaggacat 300
gaagaagtcg gaggcgatca tcccgctcca ggccggttgg aacaagggag actcgcagaa 360
gttgatgacc aacttcggca cgccccgtaa caccaccacc aagggtcaaag tggagaattt 420
ggcggaattt ccggaggaca ttttgctgaa aggacacggc gaggtgcgcc tgcagtcggg 480
taccaaccgg ttgcggtccc agaagggtt cgtcgcgttc ggtaccggac gtgacgtgtg 540
ccgtgagggg gtgaacgtga acgtgctgcc gggcgacttg gagccgcttc cggagga 597

```

&lt;210&gt; 57

&lt;211&gt; 80

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

## AKK110P1

&lt;400&gt; 57

ggcattgtgc gtctgcaagc cggtacgaac aagttcgact cgcagaaggg catgaccctt 60  
 ttcggtacgg gcccgtcgtg 80

&lt;210&gt; 58

&lt;211&gt; 513

&lt;212&gt; DNA

<213> *Globodera rostochiensis*

&lt;400&gt; 58

gaattcgcca caccgctcac atcgcggtgca aattcgccga acttaaagag aagggtggacc 60  
 gncggtctgg caagaaagt gaggacaacc cgaagtcgct gaagactggc gacgccggaa 120  
 ttgtcgaact gattccgacc aagccgatgt gtgtggaggc attcactgac tacgcaccgc 180  
 tcggccggtt tgctgttcgc gacatgaggc anactgttg cgtgggcgcg atcaaatacag 240  
 tggagaagac ggaaggcggg ggcaaatga ccaagccagc gcagaaggct ggcgcgactg 300  
 gtggcgggaa gaagacatga ccaaggggag gggcggttcc ctaaggcca accgtcgacg 360  
 aaaatgcgac caacctcttg tttatcgttg tcttattcag ttccttcac ccgtctctat 420  
 ccatattgtc gttgcgttg ataattgttt atttttgtt attgtcctgg ttgaaaaata 480  
 aatttggtca attaaaaaa aactcgtgcc gaa 513

&lt;210&gt; 59

&lt;211&gt; 393

&lt;212&gt; DNA

<213> *Globodera rostochiensis*

&lt;400&gt; 59

gaattcggtt gagcgaaaa aacatactat acaatggcaa caactgagaa gcctcaggtg 60  
 gttcaacagc ccgtgcaggc ctttggccga aagaagacag caacagccgt tgcgtttgca 120  
 aaaaggggca agggcttgat caaggtcaat gggcgctctt tggactacat gcagccggag 180  
 attctgcgca ttaagctcca ggagccaatt ctcatgtgtg ggaaggacaa atttgaggga 240  
 atcgacatac gaatccgcgt caagggcggg ggacacattg cgcaaattta tgcaattcgc 300  
 caagcactgg ccaaggcact ggtcgctttc taccagaaga atgtcgacga gcagagcaaa 360  
 aagggaactga aggagcaatt tgttgcttac gac 393

&lt;210&gt; 60

&lt;211&gt; 154

&lt;212&gt; DNA

<213> *Globodera rostochiensis*

&lt;400&gt; 60

cacgagccaa agaaattcgg tggacccggg agctcgcgct cgctaccaga atcgatccgt 60  
 taagaaataa tttttagat caaatgtttt gatgatgatc cttgtttttg ttgttgataa 120  
 aaaaaattta taaaaaaaaa ccgccgatac tgac 154

&lt;210&gt; 61

&lt;211&gt; 666

&lt;212&gt; DNA

<213> *Globodera rostochiensis*

&lt;400&gt; 61

gtattccaag tttgagcgat cagagttctt caatctatta tcaactgttt tccatcaacc 60  
 aactgtcatc atgcaaat tctgtcaagac gctcaccggc aagaccatca ctctcgagg 120  
 cgaggctagc gataccatcg agaactgtga agccaagatc caggacaagg agggcattcc 180  
 gcctgatcag cagcgtctga tcttcgccgg aaaacagctt gaagacggac gcaccttggc 240  
 gactacaac atccagaagg agtccactct ccattctctg ctgctgtctc gtggcggaat 300  
 gcaaattttc gtcaagacgc tcaccggcaa gaccatcact ttggaggctg aggccagcga 360  
 caccatcgag aacgtgaagg ccaagatcca ggacaaggag ggcattccgc ctgatcagca 420  
 gcgtctgatc ttcgccggaa aacagctcga agacgggcgc actctggccg actacaacat 480  
 ccagaaggag tccactctcc atctcgtctt gcgtcttctg ggaggagaga actgaatcgc 540  
 gggctgatgg aaagatgacg aatatgatgt ctattcgatg acttgtctct ttcgatataa 600  
 ttgattgtgt tccatttgc ggtcatcaaa tctttatgac cccctcattg ggcattggaac 660  
 gataaa 666



## AKK110P1

<210> 62  
 <211> 213  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 62  
 gaattcgttt gagaaacttt ttcaaccatt cattcaaagt tctcatcaag tgacacgggc 60  
 agcactcaac cacgggacgc gtgtactgag cgtgttggag aaggtaagt tggctctgctg 120  
 gtttgaggag acacattcgt tcgcgcaagt ggctcgaaga taccgggcag aatttggtat 180  
 ggaaccaccg cagttggacc aagtgaagaa gtt 213

<210> 63  
 <211> 488  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 63  
 agcaccggct caatcctcaa tggcacaacg acggcattct ccggcatagg agacggagtc 60  
 ggtcttggag aacaacagcc aattcccgtc gtaagcgatg cgggactgga tgcggaagaa 120  
 cagctgagaa tggccagaat gtgagccgga ggacctgaag atttatgaac gaaattttcc 180  
 agtgaagtgg accaacgctc ttcgacttta tctgctttgt gtaaagtgtg tagaatcggc 240  
 ttccaattca aaggcttttc attccccaac ttttattttt gcgcaaaaaa tttcttagga 300  
 taagcgtgaa taatttattg atttgttttt tctttctttt atctccgcct cgaagtcgca 360  
 agtgttcctt ttggcccgtt cccttttggt ttgaatgtta ttccattccc atccctcac 420  
 tttctcatat ttgtgacatt cagctgcatt gttcgactcc catttaaaag ttgagtgaag 480  
 tgcgattg 488

<210> 64  
 <211> 249  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 64  
 wccrgakbng aacahcdkdg vhwatnvcbn gschvbwagc rngtcsvddb wgnhnsswtg 60  
 gkgdyrbwnt msnwrmanrg artsstsgaa ttcccaagtt tgagagtaaa tattattagc 120  
 taaaaatggc agtcggaaaag aataagagaa tgggcaaaaa gggagccaag aagaaggctg 180  
 tcgatccgtt cacacgcaaa gaatggtacg acatcaaagc gccggcgatg ttcacacatc 240  
 gaaatssts 249

<210> 65  
 <211> 362  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 65  
 wcbcrbhdby ytsgcrsnck tbdshbcysy gcdwkmtnvk hscngdckty nyykkkvbmr 60  
 ntmsnwrman rgartsstsg tcaaccgtac tcagggaacg cgcatttcga gcgactttct 120  
 aaaaggccgc gtttacgaag tgtcactggg tgaccttaac agcactgacg ccgactttcg 180  
 aaagttccgc ctgatctgtg aagaggtaaa gggcaagatt tgcctgacca actttcacgg 240  
 aatgtcgttc actcgggaca aactgtgctc tattgtcaag aagtggcaca cgctcattga 300  
 ggcgaaatgtg gcagtgaaga ctaccgacgg ttctatgctc cgactctttt gtatcggts 360  
 ts 362

<210> 66  
 <211> 128  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 66  
 aatcaaatta agaagacgag ctatgcaaaa gcctctcagg tgcggatgat tcgtgccaaa 60  
 atggtggaga tcatgcagaa agaggctctt tccggcgatc ttgaangaaa gtagtcaaca 120  
 agcctgat 128

## AKK110P1

<210> 67  
 <211> 502  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 67  
 gaattccatt aaaaaactaa acgaacaaat ctaaagatgg ccaccgaagt ggaggaaaaat 60  
 gttcctacgg ttgacccatg ggggtgctgtg gaggaagtgg gtggtgaaga gtcgatgcag 120  
 ttggtcagcc ttgacgttac cgaggtcaaa ctgttcggaa aatggtcctt taacgatgtg 180  
 gaagtgtccg acatttcgct tgtggattat attgcggtga aggaaaaggc ggccaaatat 240  
 ctgccgcaca gcgccggccg ttaccaacag aagcgcttcc gcaaggccac ctgtccggtg 300  
 gtggaacggt tgtctttgtc aatgatgatg cacgggaggc acaacggaaa gaaactaatg 360  
 gcggtgcgca ttgtgaaaca ccccttcgag atcatcacct gctaccggag agaaccagat 420  
 ccaagtgttg gtcaatgctg tgataaacag tgggccccnc gaagatttca cacgtatcgg 480  
 acgtgcgggc actgttcgtc ga 502

<210> 68  
 <211> 519  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 68  
 gcaaaactttt atcaaaataaa aaattttatat ttgccaaaaca aattttatgaa taaaaattca 60  
 ttaatcatta aaactacatt taaaatatac tttttagaga atgtcgtcta aaatatttctt 120  
 tttctccctt tatgcattcta tctaaccaga cttggaagca atatggctaa tcaagtcaac 180  
 aatacggcag gaatacccaa actcgttatc ataccagcta accaatttaa caaatgaggc 240  
 gttgagaacc ataagagcct cggcgtcgaa aatagacgaa tgagtgtcgc caagaaagtc 300  
 ggtagaaaca acctggtcct cagtatatcc aagaatccct ttaagctttc cttccgaagc 360  
 agtcttaatt gcatttctaa tagcctcctt cgttgctggc ttctccaaac gagcagtcac 420  
 atcaacaacg aaaacgtttg ggcgtcggca cagaaaaagc catttccggg aagcttccca 480  
 tccaattcat ggattgacct ttccaacagc ctttgcagc 519

<210> 69  
 <211> 218  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 69  
 ttgattcttt attagtggac aatgacggaa gaccagaaga agttgccgat ggtgcctgag 60  
 actgttttga agcgaaggaa agttagggct gctcagcgtg cttctctact caagaataaa 120  
 ttggagaata ttaagaaggc taagggttaa acgcaagtta tctttaaacg tgctgagcaa 180  
 tacttgattg catatcgacg taagcaaaag caagagtt 218

<210> 70  
 <211> 293  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 70  
 taagaaagca gggaattttt atgtcccaga tgaacctaaa cttgcttttg ttgtgcgtat 60  
 taagggaatc aacaagggtta atttaaattt gctataaagt ttaggatggg tttagacaat 120  
 tcttctcttt taatgctttc taactttttc aaaaaagtta tgattttatc acccattaat 180  
 ctacaaattc ttttaatttat cagatccatc ctgcctctcg aaaagttctt caacttttcc 240  
 gcttgctgca aatcaacaat ggagttttca ttaaattgaa taaagctaca atc 293

<210> 71  
 <211> 422  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 71  
 aatgcaatta agactgcttc ggaaggaaaag cttaaaggga ttcttggata tactgaggac 60  
 caggttgttt ctaccgactt tcttggcgac actcattcgt ctattttcga cgccgaggcg 120  
 taagttttga ttttctaaga ttatatattaa ctttttaaat ttttcagtct tatgggtctc 180

## AKK110P1

```

aaccgcgcat ttgttaaatt ggtagctgg tatgataacg agtttgggta ttcctgccgt 240
attgttgact tgattagcca tattgcttcc aagctgggtt agatagatgc ataaagggga 300
gaaaagaata ttttagacga cattctctaa aaagtatatt ttaaattgtag ttttaattgat 360
taatgaattt ttattcataa atttgtttgg caaatataaa ttttttattt gataaaagtt 420
tg                                     422

```

<210> 72  
 <211> 374  
 <212> DNA  
 <213> *Meloidogyne incognita*

```

<400> 72
atctgagcat aaggaaactt ggcctcaagc tatagagcag accgattatg tggcaccgac 60
tgagccagtt aaactggact tcaacgttcc gcttattagt gattgggctg ctgcttctga 120
gtggcctcaa gaagaggaag ctcagggttg acctactgca ccaattgggtc agccacagcc 180
tcaacagcag caaactcaac aaggagggtga ttggaactct ggtactagtg gatgggtgaag 240
ggcaggaaaa ttgatagaaa gagaaattat tatggaataa atgtaataca tgtgtgtgtc 300
tgatttattt gttacatata caacaagttt tatittgttg tttatttaat aaaagttgtt 360
aattaaaaaa aaaa                                     374

```

<210> 73  
 <211> 120  
 <212> DNA  
 <213> *Meloidogyne incognita*

```

<400> 73
tttttttttt tttttcttca tcaatatttt gaagtgaaga accagaagta gttgcattcg 60
agctttcaaa ttttgttttt tgattactct ttaaacaaga ttcaactgat ggatctactg 120

```

<210> 74  
 <211> 369  
 <212> DNA  
 <213> *Meloidogyne incognita*

```

<400> 74
gtctaaccac tctagagcta ttcggttcgt ctgtctgttg attattagat gttgattgaa 60
cagcactagt ctctgatgta gttttcttca atctcatttt taagtgatgt agaggaagtt 120
tagaattctg attgctatcg tcttctttct cttcttttaa tggctttttc aatttatctt 180
cttctttttc ttgtccattc ttttcttcat tcttttcaaa aggtcagga aattttaatt 240
cagaccgctt ctttttaact gctgtatcta aagaaaaacc tctaggcaac gtcccagttc 300
cactcaaatt caattttgtt aaatttttgc cagatctaag tccttcttcc ttttgaacga 360
attgaactg                                     369

```

<210> 75  
 <211> 529  
 <212> DNA  
 <213> *Meloidogyne incognita*

```

<400> 75
ttttgttttt tttttttttt ttatcagaaa aaagtttaat cagaaaaaaaa aattaaaaca 60
aatctaaata aggcctctatt ctaagtttat atttttcttt tacataaacc gtcaaccctc 120
caagtttttc aatgcttgga ggttttaatt gatcctctgg taataatttg taggctagaa 180
aaaagttttg agcaaaaagg aaaaagcatca ttcttgctaa ggcttctcca gcacattgcc 240
ttttcccccac accaaaagct attagctcgt cagctttttt taatttccct tcattgtcta 300
tataacgttc aggggtcaaaa ttttggggat ttgggtatat ctttggatca aaaagaacat 360
ccgatacttg ggggtacata aatgtacctt taggcaacac aaactttcca acattcaaat 420
cttccaaggc taaatgccc aaattgaaag ggactaaatt aacgagtctt aatgtttcat 480
taacaacagc atttgtataa attaatttag gtctgtgttc caaactaat 529

```

<210> 76  
 <211> 449  
 <212> DNA  
 <213> *Meloidogyne incognita*

## AKK110P1

<400> 76  
 agtttttttt tttgaataaa agactttttt ttattaaaat ggcttcgcaa actgcaggaa 60  
 ttcaacaatt acttgcaaga gaaaagcgtg ctgcagaaaa gattaatgag gcacgtaaaa 120  
 gaaaggcaca acgacttaaa caagcaaaac aggaagcgca agctgaaatt gacaaatata 180  
 gagaggaacg tgaaaaacgt tttaaagagt ttgaacataa ttacctcggc gctagagatg 240  
 atattgctgc acaaaataag cgtgaaactg atgagacgct taatgaaatg actcgtagtg 300  
 ttgctgctaa taaacagcag gtaattgttc gtctacttca acttgtctgt gacattcgtc 360  
 cagaactgca tcacaattta caacttcaac ttaagcttaa tgaaaagcct gcctaatttg 420  
 tagttgattg attataaaaa tgaaattga 449

<210> 77  
 <211> 643  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 77  
 atttatattt gaacaaataa ttaacaaaa aagtatggct cgaggaccaaa agaagcattt 60  
 gaagcgtttg gccgctccaa agaattggat gttggacaaa ttgggtggag tttttgcccc 120  
 acgtcccatg tgcggggctc acaagcttcg tgaatcgctt cctcttattt tgtttcttcg 180  
 taatcgtcta aaatatgcac aatcttataa tgaagctagg atgatttgca aacaacgtct 240  
 cattaagttt gatggcaagg tgcgtacaga aatgcgcttt ccagctggat ttatggatgt 300  
 gggtttccatt gagaaaaact gcgaagtctt tcgtcttctc tatgatgtca aaggacgttt 360  
 cattactcat cgcatacaaa aggaagaagg tcagcttaaa ttgtgcaagg tagtaaaagca 420  
 agcgaattggg ccaaaacaag ttccttatat tgttactcat gatgcccgta ctattcgtca 480  
 tccggtacca cacatcaagg ttgacgacac tgttgctgtt gatataaaca ctggaaagg 540  
 tacagatcac attagatttg attctggtaa tgtttgatg attactgggt gtcacaacat 600  
 gggacgtgtt ggtattgggt gacatcgtga acgccaccct ggt 643

<210> 78  
 <211> 584  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 78  
 atttcctcta aaaatgaatt taaaagaaca acaaatatat ttaaattatc aattattatt 60  
 ttttattttg gctgtcagta gttttttgac aactaagggg agtgaagtaa aacaacgaga 120  
 aaataataaa ttggaatata ataaaaatga aattgagagg caaaaagagc aattaattcg 180  
 agatttgatt gcctccttaa cactgtaaag gcaatattca cgagattggc aacaatcaca 240  
 acagcaacaa aatttcatta acagttttgg cccttcccca catttatcc cctcttcagg 300  
 cattgaatgg ccccaacaac aacaaaaaat atttttggaa gaagggggaag tagaagaacc 360  
 ttttagaggaa aatgagaagg aaaaaagagc acaaaacttt gttcgtttcg gaaagagagc 420  
 acaaacattt gttcggtttg gaaaaagggg acagactttt gttcgatttg ggagagattc 480  
 aaaacatcaa cataacttgt cagatcagaa gcagttaaaa actgacaaac aataaaaaatg 540  
 atgaattatt taaaaatttt tttaatgatc ttttaattaa aatt 584

<210> 79  
 <211> 556  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 79  
 atcaagcatt aaatatgcag atttttgtaa agactctcac cggaaaaact attactctcg 60  
 aggttgaggc ttctgatacc attgagaatg ttaaggcaaa aattcaagat aaagagggtg 120  
 tcccgcctga tcaacagcgt ttgatctttg ctggtaagca acttgaagat ggacgaacct 180  
 tggctgatta taacatccaa aaggagtcta cacttcactt agttttacgt cttcgtgggtg 240  
 gaaaggttca cggttcattg gctcgtgctg gaaagggtcg tgctcaaaact cctaagggtcg 300  
 aaaagcagga acataagaaa aagaagcgcg gccgtgcttt ccgtcgcatt caataaacc 360  
 gtcgcttcac caatgttgct acttctgggg cgggacgccg tcgtggccct aactccaacg 420  
 ctgcataaga gaatggctgt atcttgatga atgtatgggt atataatcaa tttatacat 480  
 tcgactntat gaagttttct gttattcaag ataaatcttt ttgttgaaaa aaaaaccaag 540  
 tttgagatca gttact 556

<210> 80

## AKK110P1

<211> 424  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 80  
 aacattgttt taattaaaat ttacccttcc ttagcaatg acatcagaca gacttggccc 60  
 agtagttcca gatttgacag cccaagagac caacagactt gaacgaacta gttctttggg 120  
 cgatttgga attcgggatg gagttccata tccacctagg cctgcaatta ataattgttc 180  
 tccataacctg aatatgttga ctggaacgtt ttctgtacca aatgtaaatc agtacacggg 240  
 tgcaataggc ctttatcgac cagcaaatcc tgtttatact tattatagct ataaatgcta 300  
 ttttccgtat agaaattatc gaggtacac actgacggat gcttactggg acgaccgtta 360  
 ttattatttt tcgccaatat acaaacggtc aatgttccca attagattcc ggcattctga 420  
 ctac 424

<210> 81  
 <211> 89  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 81  
 attatccaca cacctattgg agctaccctt accaaggaaa atggtacgac tatgacaatc 60  
 caacanatta ccgccattc ttgaccca 89

<210> 82  
 <211> 168  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 82  
 tttttttttt taaaatttat tcattaacaa atgaccttaa cagataaaaac ttaacagtca 60  
 aaagacaaca taatttccaa ctttttcaat attatccttt ttaacgggtt gattttgcaa 120  
 ctgctccaa ttcgtccttc ttcttgatag catatgaatt gctcgaac 168

<210> 83  
 <211> 67  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 83  
 aattcatcag ccagacattc agcaattggt ttgatattac ggaaagaagc ttcacgagac 60  
 ccagtac 67

<210> 84  
 <211> 42  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 84  
 taacacgacg aagaggcgaa acatcaacag cctgacgacg aa 42

<210> 85  
 <211> 429  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 85  
 tatacgagta gaatcctccc gtggtcctcc attaataaca gcgccaacaa gtatttgaac 60  
 tggattctct ccagtcaaaa tatgtataat ttcaaaagcg tgcttcacaa tccgaacagc 120  
 catcaacttt ttaccattgt tacgtccatg catcatcatc gaacaaacca aacgttcaac 180  
 aatcggacaa tgagcctttc gaaaacgttt gatttgatat cgaccagcac tgtcgggcaa 240  
 atatttggcc gatttgctt taacagcaat ataattccact aaagaagcat cattaacttc 300  
 gatatcgctt aaagaccatt taccaaacaa tttaatttca ggaaaatcaa ttgtagtcac 360  
 ttgcatatcc ccttgtccac caggaaacac agttgcgccc caattatcat cagcgggtaa 420

accatctcc

AKK110P1

429

<210> 86  
 <211> 435  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 86  
 tttagagtttt taaaaagtac atactatttta atttttaaca aattattttg atcaatttta 60  
 aatttttcttt tcatcatttt ttaattttaaa aaacatttta acaaattaca agaacaacaa 120  
 acataattgt ctccttttta ttataaaatt taaagttaa taagttttaa aacatttctg 180  
 actggagtag gtgtacttag tgttttagaa aaggcaaaat tagttgttg gtttgaagag 240  
 acaaattctt ttgcacaagt agcgagaaga tatcgagcag aatttggaat ggaaccccca 300  
 catatggatt tagttaaaaa attacatcaa cgttttctca atactggttc tgtttctaag 360  
 ggaaatactg aacattttga agttaatcca acaatggaaa catcgacatc ctcaacagag 420  
 ggtgtagcag atccg 435

<210> 87  
 <211> 501  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 87  
 gttttttttt ttttttttta aacaaaatat cgagtcctta taagacaaaa ataaaagaca 60  
 aaagcaattt agttttatca ataaaattaa aaatagtcaa tgtctcgtt cactcattag 120  
 atttgtggcc cttaaagagg ccgtttggtt ttggttggtg tacttcagct gccttccacc 180  
 aattgttcct tagccacca atccgtaaag agtacgtcct tggcgtttca acgcatagac 240  
 gacgtccatg gctgtgaccg tctttctctt ggcgtgtacg caataagtta ccgctgcgcg 300  
 gatcacattt tcaaggaaga ctttcagaac acctcgatc tcctcgtaa tgagcccga 360  
 aatacgttt actccaccac gacgtgccaa tcgcccgaat gccggtttg tgataccttg 420  
 gatgatatca cgcaagactt ttcggtggcg cttagcgctt ccctttccaa gtccctttcc 480  
 gccttttact cgtccggaca t 501

<210> 88  
 <211> 270  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 88  
 ggaagtgtgt ttaagataaa tggatgatta gaaataaaaa tgaattgatt aaaaattacg 60  
 ttagaataat aatggaatat ataaaaataa attggatgat ttaataaaaa aaaaaaagag 120  
 agaactagtc tcgagttttt tttttttt tttttaanaa ttaacaattt atctcatttt 180  
 cctcttccat gaaaattaac aaaaagacga caacttaatc ccataattaa catcattttt 240  
 aagcttcagt cggcatgctt cgaataatgt 270

<210> 89  
 <211> 286  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 89  
 caagcggttc ccaactcaat gttgttgcca tgatactcgt gaacaccagt tctcgccaac 60  
 atagaatagt actcaatctc actgcttcta aggccttgag tattattcga aataataaca 120  
 agtttagcct ttccagaacg aagagtcttc aacgtctgct ttagcccaa acaatacttg 180  
 cccgatttgg taaccatggc gagacgagca ttgatattt ctgtggactt tttctgtttt 240  
 ccaacaacca ttgtaacgca aaattaaaat ctctttttta acaaat 286

<210> 90  
 <211> 391  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 90

## AKK110P1

```

agatatgaca tcagacagac ttggcccagt agttccagat ttgaccagcc aagagaccaa 60
tagacttgaa cgaactagtt ctttggttga ttttagcaatt cgggatggag ttccatatcc 120
tcctaggcct gcaattaaca atgttctctc atacctgaat atgttgactc gaacattttc 180
tgtaccaaat gtaaatcagt acacgggtgc aatagggtcct tatcgaccag taaatcctgt 240
ctatacttat tatagctata aatgctattt tccgtataga aactatcgag gctacacatt 300
gacggatgct tattggtacg accgttatta tcttttttcg cctatatata aacgggtcaat 360
gtttccaatt agattccggc actctgacta c                                     391

```

<210> 91  
 <211> 131  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 91
attatccaca cacctattgg agctaccctt accaaggaaa atggtatgac tatgataatc 60
caacaaatta ccgcccgttc ttcgaccac gcacacagc atcattttca agaccctatg 120
attacacatc a                                     131

```

<210> 92  
 <211> 571  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 92
ttggtgacgac aacaaaaaaa ttttatttat tttttaacaa cagaaaaata tactttttta 60
tttttaatat ttttccatga ttcaacagcc atactttcct cattttaata cttcttaaac 120
cctcaaaaaa ttcattttatt gacgaccagc agcagggtgt tgctgctggt gttgaccacc 180
acccccctgc gcttgacctt gctggtgctg tcccttcacg tcaacaggca aattgagttg 240
caaataatca accatctcct tagtctcttg atcaacacta atagttggat gttgagaagc 300
atcaagatag gaaacttctg gaacccaatt atcacgacgc tcacgctctt cttcttgcaa 360
tttaatataga attccacgaa cagggtcctt ttcgatacgc ttcatacaat gggtaataaa 420
accagcaatt tgattacgca tccgtttgct aggaataaca gcaatttcct cacaaattcg 480
tttggttcaca tgaaaatcat aagtcaagcg tgtataatat ttgtcaataa taacacgaga 540
tgctttcttg acagttttga gagaaccgat t                                     571

```

<210> 93  
 <211> 671  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 93
tttgagaatt taacttttct aacaaaaact tttatttttg tctttgatgt ctactcaagt 60
accaatacgc gtgctggtta ctggagcagc tgggtcagatt gggtattctt tgggtattca 120
aattgcaaaag ggtgatgttt ttggaaagga aacgcccatt gttctggtta tgggtgatat 180
tcctccaatg gccgaagtgc ttaaaggagt ggaacttgaa ctttacgatt gtgcttggc 240
gaatcttata gctgtcgagc cagtcacgac tgaagaggca gcgttcaaag acattgatta 300
tgcttttctt gttggtgcaa tgcctcgaaa ggaaggaaat gaacgaaagg atttacttgc 360
tgctaattgtg aaaatattta aatcgcaagg attggtctta gcaaaatatt caaagccaac 420
tgtttaagggt ctggttgttg gaaatccagc aaatacaaat gcttttattt gtgcaaaata 480
cgcagcagat aaaattccag caaagaatgt cagcgctatg actcgtcttg accataaccg 540
tgcaattgcc caaatagctg ctcgttgttg ggttgactgt ggatctgtga agaaagttat 600
aatttgggga aatcattcaa gtacccaatt tcctgatgtt aaacatgcta aagtaattaa 660
agggtggcacg g                                     671

```

<210> 94  
 <211> 289  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 94
ggctgtaaaat gatgtgccgt ggatacagaa tgaatttatt tcgaccgtcc aaaagcgcg 60
agctgttatt atcgaaaaac gcaaaactgtc cagcgcaatg tcggcagcaa aggcggcatg 120
tgatcacatt catgattggc acttttggaa aaaagatggc gattgggttt ctatggccgt 180
tccttcgatg ggttcttatg gaattccgga aggtttgatc ttctcatttc caattacaat 240

```

AKK110P1

tgatgcanaa acgcgtgact ggaaaattgt acaaagatta gaactcgat

289

<210> 95  
 <211> 262  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 95  
 aatttaactt ttctaacc aaacttttatt ttgtctttg atgtctactc aagtaccgat 60  
 acgcgtgctg gttactggag cagctgggtca gattgggtat tctttgggta ttcaaattgc 120  
 aaaggagat gttttcggga aagaaacgcc catcggtctg gtaatgttgg atattcctcc 180  
 aatggccgaa gtgcttaaag gagtggaact tgaactttac gattgtgcct tggcaaactc 240  
 tatagctgct gagccagtca cg 262

<210> 96  
 <211> 323  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 96  
 aagacattga ctatgctttt ctgtgttggt caatgcctcg aaaagaagga atggaacgaa 60  
 aggatttact tgctgctaata gtaaaaaatat ttaaatcgca aggactggct ctagcgaat 120  
 attcaaagcc aactgttaag gttctgggtg ttggaatcc agcagatata aatgctttta 180  
 ttgtgcgaaa atatgcagca gaaaaaattc cgacaaagaa ttccagcgct atgactcgct 240  
 ttgaccataa ccgtgcaatt gcccaaatag ctgctcggtg tgtggttgac tgtgggtctg 300  
 tcaagatagt tataatgtgg gga 323

<210> 97  
 <211> 717  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 97  
 aatattttta acaaacgatg taacagaaaa acaagtttt tttaacaaat tttcttgaac 60  
 cttatttttt ttcaaaaacat ttttttattt aaattttaaac ctctcttcat ttctcttaaa 120  
 cactttcctg aactggaggt tcataagcat ctggacgact ttcaataact tctecacttg 180  
 ctgtagttat agcaacttgt ccaccaccac ttccagcacc ctctccatgc atatccaaaa 240  
 gttttccaag ttcaaatttt gggtttttca aaatttttac ttttcgaata taaacgtctt 300  
 gaagtggata gaaataagaa caagactttt caatgtcttt tccaatagaa tcaggaaatta 360  
 atttgcgtac aacttcttta agatcgcatg aagaaacctc gcgtagaata atctcaacca 420  
 tcctagcacg aatttgacgc acttgagacg attttgcata actagtcttt ttcacttggg 480  
 ttggagcttt ctttgtgaag ccaatacaga acaatcgaag caaataacca tcagtgtgtt 540  
 tgacagcaac atttgcctca attaaagtat gccacttttt gacaatagaa caaagcttgt 600  
 ctcgagtaaa agtcattcca tggaaattgg tcaaacaaac ttgccttga acctcttcac 660  
 aaataagtcg aaatttgcca aagtcagctt cggtgtgtgt cagatcacca agagaaa 717

<210> 98  
 <211> 758  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 98  
 gacaagttta accttgtgtg actttatcta tattcttgtc taaataattc taacaaattg 60  
 taacaacaaa caaaaatggg cgagcaagac aaaaagaaag ctggcggcgg cgatggtggc 120  
 aaaaagaagg atggcttcga tgccaaaaag ttgacgattg atttggcttc tggaggaact 180  
 gccgctgcgg ttcttaagac ggctgtggcg cctattgaac gtgtcaagtt gttgctacag 240  
 gttcaagacg ctctcagca catcgctgcc gataaacgct ataaaggaat aattgatgtg 300  
 ctgtttcgtg tgcccaaaaga acagggagtc ctgtcttttt ggctgtggtaa ttgggctaac 360  
 gtgatccggt actttccaac gcaagctctc aactttgcgt tcaaggacac ttacaagagg 420  
 atcttcatgg aagggtgttg caagaacaaa cagtttggca aattcttttt gatgatgctg 480  
 tttatgagca aaaatttcct tgggttggat agacctaaca gttgaagagt atcttgccct 540  
 ctgtgatagc tatacaacac tctcttcaat tggagattca atgttgagtg gagatgctga 600  
 tagtaatcct ctgttacaat cacttaacaa ctcaatcaat tccaatgcca ctgctcagaa 660  
 ttataactcc tcaacaattg gccgaagcta aaactacgt ttcaacatgc tacagctact 720



AKK110P1

tcaaaatcga aacagattgt tttaaactgt tgaattt 758

<210> 99  
 <211> 154  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 99  
 ttgagttcgt tggcacattt gttgtgttac aaaacgaaaa ttattgggaa cgggtttcag 60  
 tgcctattct cgcaggttat tggcattca cacatttgta ccaataacaa cgttaccgtt 120  
 tataatcaaa ctgttcctca aagttatgcc catt 154

<210> 100  
 <211> 125  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 100  
 ttcagaatac tcaaggctctt atattcgttg ttgatagtaa cgacaaagag cgtattgttg 60  
 aagctcgtga ggaattgatg cgtatgttgt ctgaagacga acttcgcgat tctgtactcc 120  
 tcgta 125

<210> 101  
 <211> 219  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 101  
 cttgccgaat gctatgaacg ctgctgaact tacagacaaa cttggacttc acacgctgag 60  
 aaatcgtaac tggatatatcc aggtacttg tgccacttca ggagatggtt tgtatgaagg 120  
 ttggactgg ttgagtaacc aattgaagaa tcaagggttaa atgagctctaa ataaaaatgg 180  
 agaggggaaa gaggagaggt taatttttta aggaaaaaa 219

<210> 102  
 <211> 473  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 102  
 gttttttttt ttttttttta aattccaagt tttcttccaa atgagagaat agggagaatg 60  
 atgggggaaa aaataggagc aagccaaaaa gccaaaaaaa aattttttt ttaaagtatt 120  
 tttgtaaatg tgtgaaaagg tgtgtgtcaa ttgtagagtc aaatgtcgtt gccttccttc 180  
 cactaaaatt tctctttcct ttcttttctc ttctaaaatt ccttcaaagt cgatccaacg 240  
 aaatttcagc ctctcttgga tattccaact cccaaatcag cttcaaagt ttgcctttaa 300  
 cgtcacgagg agtaccaaat ccagtcatca acttttgaga gtctccctta ttccaaccgg 360  
 cctgggatgg aattatcgtt tctgacttct tcatatcttc atatggaagt tcgccagact 420  
 ccgcctcgta tgttgtgttc cttggcggtc caaaacctgt catgcccgtc tgc 473

<210> 103  
 <211> 114  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 103  
 ttggaccgtt aggattgtcg ccaaagaaaa ttggagaaga cattgcaaag gcaacacaag 60  
 actggaaagg cttaaagggt acttgcaaat tgactatcca aaaccgaatt gcc 114

<210> 104  
 <211> 255  
 <212> DNA  
 <213> Meloidogyne incognita

## AKK110P1

<400> 104  
 ccgcttctcg aattgtgaag gaattgaagg aacctcaccg agaccgcaaa aaagtcaaac 60  
 acgtaaaaca cagtggaaat ttgacgatcg agcaaatat caacattgca cggcaaatgc 120  
 gacctcgttc aatggcgaaa aaaattggaa gggactgtta aggaaattct tggcactgca 180  
 caatctgttg ggtgtactgt tgatggacaa catccacatg atattgttga tgcaatccga 240  
 agtgggaaaa ttgaa 255

<210> 105  
 <211> 571  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 105  
 tttttttttt tttttttttt tgtcaacaat aaattttactc agaaaaatca ttttaacaatt 60  
 taacacacat ttttaattcc ttaatactcc aaaaaacttc tcttctttat tccctcttat 120  
 tctcccaatt catttaaagt ttcagttttg tgcggcgcca atgacgacgt ttgcatatat 180  
 agcgtatacg actgccagtt ttcattcgaa cccattgctg cagcggtcga ttttgtttag 240  
 cagccttagc cagcttgctc ttgataataa acgtttttgtg tgcagccatt aaattgttga 300  
 ctttatccaa aattgttttt ttgaaggcaa taaacaaatt taatttttct gctcaacaag 360  
 tccatagcag ctcatctggt caacaatctc cctcatgctc ctcagctctc agcgtcttct 420  
 cttatgaatg tcaaaaacag cagcaacaac ccccagcaga acctgttggg ctttctttgg 480  
 aagttcatca atctgtgcat tcaacaacaa ccttccatc tccatgttct ttattacccc 540  
 ctccctcttc tttacatcct ataatcatc g 571

<210> 106  
 <211> 235  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 106  
 tgctttattt tcaattcttc aacccaaaaat taaatcttcc cttattttta ttacaattcc 60  
 aattttagca gcattagccc caactacttt agctgctaataaaaattgttt atgaggatgg 120  
 agatagtgat ggacttgata tggctaaaag tattttaaat tgaataaagg aaaaagaagc 180  
 attttaaaga aaatttagatg gaaatgctga agaaagaaaa aaattattta ttttt 235

<210> 107  
 <211> 702  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 107  
 ttttttcaaa aaataattcg aattttgttc ttttttatTT tgctacaaat aaaattttaa 60  
 ttgaaaaaaa aaaaaaaaaa aaaaaaaaaa tgcagaagaa atccttgccg aaattgacgg 120  
 ctctcaaatt gaggagtatc aacgtttctt cgatatgttt gaccgtggaa agaattggcta 180  
 tattatggct actcaaattg gggtaattat gaatgctatg gaacaagatt ttgatgaaaa 240  
 aactcttcgg aaattaatcc gaaaattcga cgcagacggc agcggcaaaa tcgaattcga 300  
 cgaattctgc gctttgggtat acactgtggc gaatactgta gacaaggaca ctttgcggaa 360  
 agaattgaga gaagcttttc gtctctttga caaagagggc aatgggtaca tctctcgtcc 420  
 aacactcaaa ggattacttc acgaaatcgc cccagacctc agcgataaag acctggatgc 480  
 cgcagtagac gagatcgacg aagacggaag cggaaaaatt gaatttgaag aattttggga 540  
 gttaatggct ggagagactg attgaaattt taattagaat gactagaaaa ttgactaaaa 600  
 tattttgccca ttaaattttg gaaagtgccca aaaattgcct ttctgagaat ttttattttt 660  
 aacgtctaaa taatgaataa aatggatata aaaaaaaaaa aa 702

<210> 108  
 <211> 423  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 108  
 aaaattaaaa taaaagacaa acaataaaat ataaattaaa taaataatat ttaaataaac 60  
 acacaaataa actctccaaa cataattttt ttaaaatttta ataactttt gtcccatTTg 120  
 agaaaagaaa tgccaaagga gatgaagaac ttgttgaaga aaaaagttca aaaatatcaa 180  
 ctccctcatt tgcgtcaca ttttctttca ttattccatt tgttgtaagc tcagtaactg 240

## AKK110P1

```

cccccaattgt tgtttagtc catggagaga aagcactttc cccattcgaa aatgttgaac 300
caaatttggtc aaattgttgc tgtttagtc ctcgaagttc gttagaaaca gaacgaaata 360
aattatgagg ttgttgttgc tcctgacgtt tttgattgtc tggagctggg tgaggatcac 420
caa

```

<210> 109  
 <211> 994  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 109
ttttattttt tatttgaaaa taatcatcac attataatta atgggaaaaa gacaaaaaat 60
tagaacaggt gctggcgatc ttgtcacaac ccctggacct cttcataaac aaattgaaag 120
gtcaaaaacta gccaaagccga aattcaagcc tttaaaacgt tcaagagaag agcaaaaaga 180
tgaaattgaa cttgtcgatc catcgtaaa gggcaaaatt attattaaag caaacaataa 240
attggaaaaa gatgttgtgt tcaatgagga tggagaatct gataattctg aagaaaattga 300
agaagaagaa gaagacggca atgaaaagtt ggatgttgat caattagtat caaacattt 360
ggaagattta gatgaactaa aattggatga tggcggtgaa aatgtgcgaa agataataac 420
gaaattcaga taaaaataac aaagaaaagt ttataaataa agctgagttt gccgatatcg 480
acccaaaaat tgttgatctt ttacagaaa ttggtcaagt tttaaagaaa tatagaagt 540
gacgtattcc caaagctttt aaagttattc caactttggt tgattgggag aaaattatcg 600
aattaactcg cccagatgat tggtcggcag ctgcaatggt acatgctacc aaaattattg 660
cttcaactgc taccctact caatgccaaa gggtttataa ttgattttg ttgccacgta 720
ttcgagatga tattgacgga ttaaaaaatt acatttccat atgtatcaat gcttatttaa 780
agcattgttc aaaccagctg catttttcaa aggaatcctt ttgccgctt gcaaatcgaa 840
caatttttct cttcgagaag ctgttgttct tgcttctatg cttcgtaaag cttccatccc 900
tcaattacac gcggccgcag catgtttgag tatttcttgt ttagaatata cttcttcaag 960
ggcttatatc cttcaagcat tgatagaaaa gaat

```

<210> 110  
 <211> 476  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 110
tttaaacact taaaaatacc ttcaaattta ttttagaacc tttttgccat taaaaaaaat 60
tttatttcga aaaaatggct gagaatatag aagaatcct tgccgaaatt gacggctctc 120
aaattgagga gtatcaacgt ttcttcgata tgtttgaccg tggaaagaat ggcttatatta 180
tggccactca aattggggtg attatgaatg ctatggaaca agattttgat gaaaaaactc 240
ttcgaaaatt aatccgaaaa ttcgacgcag acggcagcgg caaaatcgaa ttcgacgaat 300
tctgcgcctt ggtatacact gtggcgaata ctgtagataa ggacactttg cggaagaat 360
tgagagaagc ttttcgtctc ttcgacaagg agggtaatgg ttacatctct cgtccaacac 420
tcaaaggatt actccacgaa atcgccccag acctcagcga taaagacttg gatgcc 476

```

<210> 111  
 <211> 189  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 111
cgaagacgga agcggaaaaa ttgaatttga agaattttgg gaattaatgg ctggagagac 60
tgattgaaat ttaattaga gatgaataaa aaattaacta aaatattttg ccataaaaat 120
ttggaaagt ccaaaaattg cttttttgag aatttttatt ttaacgtct aaataatgaa 180
taaatggat

```

<210> 112  
 <211> 164  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 112
ttgaggaaat ttaatttttt aaacaaatat aataattacc aaacaacaaa aaagaatccc 60
aaaaacaaca tttttaaatc aaatgacaga catatatttg caataacgat gtgtggattt 120
tctttttttt taaataatta acatcttaag cctgctattt cttc

```

## AKK110P1

<210> 113  
 <211> 539  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 113  
 cagctttctg cgcagatttg gtaacctttc caccagcttc gacctttctg acggccttga 60  
 taacaccaac agccacagtt tgacgcatgt caccgaacggc gaagcgctca agaggagcgt 120  
 agtcagtaaa agcctcaaca cacattggct tggttggaat taagtcgaca ataccagcat 180  
 ctccagtcctt caaagccttt ggattgtctt caacctttctt tccagttcga cggtcgacct 240  
 tctctttaag ctccagcgaac ttgcaagcaa tgtgagcagt gtgacagtca agaacaggcg 300  
 tgtagccagc agcaatctgc ccaggatggg tcatgatgat aacctgagca gtgaattgct 360  
 tgggtctcctt tgctgggtca ttcataagagt cagaagtgc tgaaccacgt cggatgtcct 420  
 tgacagagat gttcttaacg ttaaattccaa cattgtcttc aggaacagct tcagggagag 480  
 actcgtgggtg catctcaaca gatttaactt cagtagaagt tccttcagga gcaaaagga 539

<210> 114  
 <211> 314  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 114  
 gtttttaatt ttagaaaatg tctacagaâa cagaaaagga tttagaacgt tgggaggatg 60  
 tccgtcgatt tactgagatt ggttcttcta aatttgccca tcccgtttt gttccaagcc 120  
 cggagaatct tgaaagagta aggaaatgct cagttttggt tgttggtgct ggtgngcttg 180  
 gatgtgaaat ttgaaaaat ttggccttat caggatttca aaatattgaa gttattgata 240  
 tggacacaat tgacctttca aatctcaaca gacagttttt gtttcgtgaa cacgatgttg 300  
 gcttatacaa agca 314

<210> 115  
 <211> 200  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 115  
 ttcgaagacg tggttaaagga tgtcgtctta ctgcacataa ttgtaaaata caagataaag 60  
 gacttgactt ttatgggcaa ttttcaatta taatttgtgg actagattct attgatgctc 120  
 gaagatgggt aaacgccaca gtgtgttctt tggtcgaatt tgacgaagaa aacaagccac 180  
 ggccaggcac aattattcca 200

<210> 116  
 <211> 471  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 116  
 tttggtcgaa aaaagactgc tactgctgtg gcatattcca aaaagggaaa aggattaatc 60  
 aagggcaatg gccgtccttt agaatttttg caacctgaaa ttcttcgtat taagctacaa 120  
 gagccattgt tgattgtagg aaaggacaaa tttgctggaa tggatattcg catccgtgtc 180  
 aaaggtgggt gtcattgtgc acaaatttat gcaattcgac agtcaattgc taaagttttg 240  
 gtggcctatt accagaaaaa cgtggatgag caaagcaaga aagaattgaa ggatcaactt 300  
 gttgcttatg atcgtaattt gcttgttgcc gatccgagac gtcacgagcc aaagaagttt 360  
 ggaggacctg gtgctcgtgc tctgtatcag aaatcttatc gttaagaagt atgaaattat 420  
 aaaattgtgt gttacgaatt aattgttatt ttgttgggat aaatntgaat a 471

<210> 117  
 <211> 593  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 117  
 gaattcaaaa aatattaaaa ttgtttaata taatttctaa aatgaagcca aaggttggaa 60

## AKK110P1

```

ttaacggatt tggacgtatt ggacgtcttg ccctgcgtgc agcggtcgag aaggatactg 120
tccaagttgt ggctgtcaat gacccgttca ttgatcttga ctatatggtc tatatgttta 180
actatgattc caccacacga cgcttttaag gaaagattca agcaagcaat ggaaatttgg 240
tagttgagaa ggagggaaaag tctactcata ctatcaaagt ttcaacttc aaagaacctg 300
aaaagattga ctgggcagggt tctggtgctg attttgttat tgagtcgact ggagttttta 360
ctactaccga gaaagcttct gctcacttga agggcggagc caagaaagt gttatctccg 420
ctccatctgc tgatgctcca atgtttgtgg ttggtgttaa tgaggacaaa tatgatcctt 480
ccaagcatca tatcattagt aatgcttcct gcactactaa ttgtcttgct cctcttgcca 540
aggttataaa tgacgagttt ggcataattg aaagttgaat gactactgga cac 593

```

&lt;210&gt; 118

&lt;211&gt; 576

&lt;212&gt; DNA

<213> *Meloidogyne incognita*

&lt;400&gt; 118

```

gaattccgag tttttttttt ttttttttaa aacaaaaatt aaaagattta tgcctatcct 60
ttgccagcca tttgcccgcc atttttttgt gcacaataaa tttttttgta atttttgggg 120
tgagggggaa gtaaaatgaa agaagggaga gagatatgaa ttggaggttt ttttgtaaaa 180
ataaattttt ttttcttgaa aattcttccc gtttctgagc ttttctgct ttttcaatt 240
ttcgtttgtc gaaatactaa actttacaat ttggttaggt tctatttgtg aaacataaat 300
atctccatta tcgctgattg caagggcatg ggcgttttcg agaccctttg caaagctatt 360
agcccttctt gtgttcatat ccattacgaa aacttgggat tctaattgac tgccttgatc 420
ttgattggtg acgccgacga ggaagtgttc tttctctcgg atagcaaaga ctgcaccaat 480
attttcagcc tttgtgaaga aagtgcctgt ggggacgtaa gcacgtctat gttggtgttg 540
agcgccttct aatccagcag aaaagcattg aatacg 576

```

&lt;210&gt; 119

&lt;211&gt; 559

&lt;212&gt; DNA

<213> *Meloidogyne incognita*

&lt;400&gt; 119

```

acgcagagta agttgagatc ttcaataagg gttagagagt gtggtacgag gaattctcca 60
tttttgggtg tttcactgga gtcaggcttc ccaaattgac tgagcaattt cccatccttg 120
tcaaaacttca ttattcggtt attacagtaa ccacttgcca cgaaaaactc tcctgtactg 180
gcaatagcaa cgtctgtagg tttgcaaaaa tgtttgcatt ctgtcccttg aacaagcttt 240
tcgccccaac tcataattaa tttaaaatcc ttgtcaagtt tgtggacttg atgacttcca 300
acgtcagtaa cccaactatt gccgtgggca tcgattgtta gtccatgagg catgtaaaac 360
atgctttttc cgtattcttc caagactgcc cctgattccg tgtctataac agcaattggt 420
gtgtttgaaa tgatgcccag ggatctgttt aggtggttgt tctcatcaaa cgaaaattca 480
tcccaaactc tgcagatcg gtgaaaaaga acaagtcgat tcaatggatc caatgcaata 540
cccggagctt gcccaatat 559

```

&lt;210&gt; 120

&lt;211&gt; 366

&lt;212&gt; DNA

<213> *Meloidogyne incognita*

&lt;400&gt; 120

```

tttaagaatt ttttaaaaat taaaacttgg actagatttt aataaaatgt cagctccacg 60
tagtgttgct agcgggtgtg gtgctgtgtt tatgaataag caagcaagta aatacaatga 120
agttgaagga gaactccttc ttaattggat taagaaagt acaggcgaaa atattgtctat 180
aaacggaact agggaaaatt ttgtgaaaca attgaaagat ggaactctgc tctgcaaat 240
tgctaacaaa atttggtccaa attcaatcac aaaggcacag gcaaaaccga acagcacatt 300
ccaatatatg agcaatttgg agctgttctt aacattttatt tcaagccaag gagtccctag 360
ggagga 366

```

&lt;210&gt; 121

&lt;211&gt; 661

&lt;212&gt; DNA

<213> *Meloidogyne incognita*

&lt;400&gt; 121

## AKK110P1

```

ttagttgaat ctcgtgacct ctactctgtt tgtatgacat taaattctct tggccgcatt 60
ttggaacgtc aaggaaaaac tcatccagag caggttaagt cgtcagaaat tcttaatttg 120
ggtactggag accaagtgcg ctttcgtgtt taaagatggg aaattgaaag aatttttggt 180
aaacataata aaaagacatt ttatggcaat aaaaaaatgt caaaaaagct tgtcttttaa 240
atattttggc aaaacatttt actttcacaa aattttaaaa taaatttatg aagattgttc 300
cgtcactttc atcatttccg atcgaccttt gttgttttct aagttcgttg gccaaagaaa 360
ggatatgtaa aattgaatta tgaataaaaa taaatcactc aatcagaggc attgttagtc 420
tctcacttcc tcctctttac ccattggcta accagcttta aggattttt ccataagttc 480
aaggtgtacg taaatcgaat accgactgtg gtatcttaat ttttccatga aattctccaa 540
taaaaaaaa tttttttat tttttttcca taatgctatc tataattttt gcttttaatc 600
ttttttggct atcaggcttt aaaatagtaa atatacttat attaatattt tatttccttt 660
a

```

&lt;210&gt; 122

&lt;211&gt; 173

&lt;212&gt; DNA

<213> *Meloidogyne incognita*

&lt;400&gt; 122

```

ggagagtttt tcgtggcaga tggttactgt aatagtcgaa taatgaagtt tgacaaggat 60
gggaaattgc tcagtcaatt tgggaagcct gactccagtg aaacacccaa aaatggagaa 120
ttccttgtac cacactctct aaccctcatt gaagatctca acttactttg tgt 173

```

&lt;210&gt; 123

&lt;211&gt; 584

&lt;212&gt; DNA

<213> *Meloidogyne incognita*

&lt;400&gt; 123

```

cgcattcaat gcttttctgc tggattagaa ggcgctcaac accaaccatag acgtgcttac 60
gtccccacag gcactttctt cacaaaggct gaaaatattg ggcgagtctt tgctatccga 120
gagaaagaac acttcctcgt cggcgtcacc aatcaagatc agggcagtca attagaatcc 180
caagttttcg taatggatat gaacacagga agggctaata gctttgctaa gggctctagaa 240
aacgcccatt cccttgcaat cagcgataat ggagatattt atgtttcaca aatagaaccc 300
aaccaaattg taaaatttag ttttctgaca aacgaaaatt gagaaaaaaa aaaaaaaagc 360
tcagaaacgg gaagaatttt caagaaaaaa tttttttacc aaacaaaaaa cctccaattc 420
atatctctcc ctcttttcat ttttcttccc ctttctcccc aaaaattaca aaaaatttta 480
ttgtgcacaa aaaaatgggc gggcgggcga atggctgggc aaaggatggc gataaatctt 540
ttaatttttg aaaaaaaaaa aaagaattcg aattatatgg ccta 584

```

&lt;210&gt; 124

&lt;211&gt; 650

&lt;212&gt; DNA

<213> *Meloidogyne incognita*

&lt;400&gt; 124

```

gtttaagaca attaaaacgt ttattttcta caatcaaac aaatatggct gttcctcccg 60
atgttatcga gaagatcgag gctgggtaca aaaagttgca ggaggcaccg gagtgcaagt 120
ctcttctcaa gaagtacttc acgaaggaag ttatggacca gtgtaaaggg ctcaaaacta 180
agcttggtgc gaacttgctt gatgtgatcc actctggagt tgcgaatctc gatagcggg 240
ttggtgttta tgcgcctgat gctgagtctt acactctctt caaacgcgtt tttgaccgga 300
ttattcagga ttaccacaat ggatttgac ctgaccagaa gcagccgcaa actgacttgg 360
gtgagggaag gactcagctt ttgcctgatc tggatcctga gggtaaatc atcaactcga 420
ctcgtgttcg atgtgggcgt tctcttcagg gatatccgtt caatccgtgc ttgactaaag 480
agaattatac ggaaatgcat gacaaagtta aaggggtttt tgagcagctt aagtctgatg 540
ctgagcttgg tggcacctat tatcctttgg agggaatgac caaagaggtt caaactcaat 600
tgatcaagga tcacttctc ttcaagaag gagaccgctt tttgcaagct 650

```

&lt;210&gt; 125

&lt;211&gt; 1013

&lt;212&gt; DNA

<213> *Meloidogyne incognita*

&lt;400&gt; 125

## AKK110P1

```

tttttttttt tttgatgttt ctaatttttg tgggcaatat ttaattattat ttttaattat 60
taaattttct tctttatttt ttaaaaaatt atttcttaaa tttattcttc tcctcttcgt 120
gttttgaaatc aaataatttaa attttaaaatt atttaaacag ctacacgagg cctcagccctc 180
ccccgttgca ttcaaattgg tcggcacggg tggcgatgat aattttattt tttaggtaat 240
tttgggtgaga aaatatTTTT aaaggtaata atgtcctttt ggacaattaa aaaaaaactc 300
gaggagagag tgaatatttt tacaatttat ttgaagagca gccagcctat tgttatcaac 360
aaaaaacctt caaaatgccg gaaaatgatt atgatgagga ggaggcgcca aacgccacga 420
tggaacaaca ggtagcttca ggtggacagc caaaacgctg ttggaaaatg gacattatcc 480
cagctgcgcc agactgatgg tataattcca tcccaggccg gttggaacaa gggagactcc 540
caaaagtga tgaccaattt tggtagctca cgtaacacaa caacaaaat tcgtgctgaa 600
tgcttgcctg atgtgcctga agaaattgct cttaaaagtc acggtgaagt acgcctccaa 660
tccggtacta accgttttgc ttcgcagaag ggaatgggtg gatttggtag tggacgtgac 720
ttatgcagag aaggagtggt tgtgagtcac gaccagccg atttatagcc cctcccagaa 780
gagataatcc gtgctagcga tggaaattgt cgtctccaat ccggtacca caaattcgac 840
tcccaaaagg gaatgggtcag cttcggtaca aaccgacgcg aaactacaag aatgaaagac 900
accaaaccatc cggaatacaa ccacgaagtt aacattgacc aaagcgaat tcctttgcaa 960
tctggtacaa acaaatcgc atcccaaaag ggaatgacca gcttcggtac aaa 1013

```

<210> 126  
 <211> 80  
 <212> DNA  
 <213> *Meloidogyne incognita*

<400> 126  
 tgttggacac tgctcaccca gaatacagtc acgaaagcag catcgatcaa acgagcattc 60  
 cttaccaaat gggatcaaat 80

<210> 127  
 <211> 585  
 <212> DNA  
 <213> *Meloidogyne incognita*

<400> 127  
 aggggaatgac ttgcttttga cagccacgtt gggaggtgct tgacccgagc attagctacc 60  
 agaaccgtaa atcacaagga atgggtccgtc tccaatccgg aacaaaccgg gtcgcctcgc 120  
 aagcgggcat gacaggtttt ggaactccaa ggaacacaa atacgaggcg gtagctggtcg 180  
 aacttcata cgaagatatg aagaagtcag aaacgataat tccatcccag gccggttggc 240  
 ataagggaga ctctcaaaag ttgatgactg gatttggtag tcctcgtgac gttaaaggca 300  
 aacatttgaa gcgtatttgg gagttggaat acccagagga ggctgaaatt tcgttggatc 360  
 gactttaaag gaattttaga agagaagaaa gaaaagagaa atttagtga aggaaggcaa 420  
 cgacatttga ctctacaatt gacacacacc ttttcacaca tttacaaaat acattaaaaa 480  
 aaaatttttt ttggcttttt ggcttgctcc tattttttcc cccatcatt ctccctattc 540  
 tctcatttgg atgcaaaactg gaattttaaa aaaaaaaaaa aaaaa 585

<210> 128  
 <211> 287  
 <212> DNA  
 <213> *Meloidogyne incognita*

<400> 128  
 catctggaga aacgttgagg caatacatcg ttattggccg taaacttctt acagagaatg 60  
 agccaaatcc aaaactttac aaaatgcaaa tttttgccag taatcatggt gttgctaaat 120  
 cgctgttctg gtactttact agtatgttgc gtcgtgttaa gaagactaac ggagagattg 180  
 tttcgtgtca ggaggttttt gaaaagaaga taggctctgt aaagaattat ggaatttggc 240  
 ttcgttatga ctctgaacc ggtcatcaca acatgtaccg tgaatac 287

<210> 129  
 <211> 175  
 <212> DNA  
 <213> *Meloidogyne incognita*

<400> 129  
 gctgtcactc aggccttatcg cgacatgggt gctcgtcatc gtgctcaagc cgatcgaatc 60  
 caaataatca aggttcaacc gatcaaggct gccgattgca aacgtactgg agttaaacag 120

## AKK110P1

ttccacaact cttcaatcaa gtttcctttg ccgcatcgtg tgaatgacaa acgtc 175

<210> 130  
 <211> 599  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 130  
 acttttgttt ataatcacat ttgcattact ttccgtccat ccttctttga gacagaat 60  
 aaagggtcac cttctaagta aggattgtag cggctgtatg attgatgttg cttttgttgg 120  
 ggagcaatag aacgcttgcg tcgccgaggc tcctcagccc tagtaacgtg aaatttcttt 180  
 gcaatcatcg atttgtgtag tccatttttg gctaagacct gttctaagtc ttgttcatat 240  
 tgttcagaat tgctttttga ttgacagtta aacatgtgtt cttggtcaca aaggcattgc 300  
 tgattggcct ggtagctacg cgagaaatcg gcggtgttat caaactcctc caaacatcca 360  
 tctcgactgg agtatccac agggcagggg tttggagggg cacaatatgc tggcaaaaca 420  
 ttgtcactct taatctcttg gcggtgtgaa aattcagatt ctggatggag ttgttggctt 480  
 ccttcaccgg caccctctgt cataaattta tgtccaaacg caatggggccc ggaagcactt 540  
 tcaatgtcac gagaaatcaa gtcgattaat tgtgaatgcg gaaatatagg ctccccaga 599

<210> 131  
 <211> 466  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 131  
 gaagattgga tttattggcg ctggaaagat ggcacaggca ttggccagag gactaataaa 60  
 ttctggacgt tatccttcac aaaatttgat ggctagttgc cctaagactg atgtctcttt 120  
 attggaggat tgcaagaggc ttgggagtaa tacagcacat gataatgcac aagttgctcg 180  
 tgaaaatgat gtggtgatta tagcaggtaa accaactatt gtgtctaaag ttgcttcgga 240  
 aattgcacca gccatccgcc gagatcatgt acttatttct atagcattgg gcatcaccat 300  
 acgctacatt gagcagtaat tgacttcaga atcccgaatt gttcgtgtaa tgccagatac 360  
 tcctgtagggt ggtaggagca ggctgctgca gccatatatc attgggatca gcattgtcag 420  
 gatagggtgat gcccagatag ttcaagatct tctgataacg ctgggg 466

<210> 132  
 <211> 266  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 132  
 atgaaattcg agttctttgc atcaaggccc gtgaaatttt tctttcgcaa cctattttgc 60  
 tggaattgga agcgccgttg aagatttgtg gcgatattca cgggtcaatac aacgaccttt 120  
 tgcggctttt tgaatatgga ggttttccgc ctgaagcgaa ttattttatt ttgggtgatt 180  
 atgtggatag aggaaagcag agcttggaga cgatttgttt gctgttggcc tacaagatca 240  
 aatcccccgga aaattctttt tgccta 266

<210> 133  
 <211> 308  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 133  
 tctatcaacc gaatatatgg attttacgat gaatgcaaac gcagattttc tataaaattg 60  
 tggaaaacat ttactgattg cttcaattgt ctgccaatg ctgctgtgat cgatgagaaa 120  
 atattttgtt gccatggagg tttgtcacca gatttgcaga atatggagca aattcgaaga 180  
 attatgcgac cgacggatgt gccagataca ggtcttctct gcgaccttct atggtctgat 240  
 ccagaccaag atgtccaagg attgggagaa aatgatcgtg gggctctctt cacttttggg 300  
 ccagatgt 308

<210> 134  
 <211> 335  
 <212> DNA  
 <213> Meloidogyne incognita



## AKK110P1

<400> 134  
 taaatttagt ttcttttctt ccatctcttt ttatgttttg aaagagtgtg ccaaaacaaa 60  
 tggccgcccg tgatggaaga agcaggcaaa attatttaca agaacattca attcctcaac 120  
 tttttgaggg tttaatgact ggacttatat acaatcaacc aatcgatcct attcaatttt 180  
 tggagaatgc aatagctaaa cttcgaaaaa atcctgatct tccattaaag tgggatactt 240  
 ttataagtgt ttcgcctcaa caacagcaac aacaacagac gagaatgaat actggagaaa 300  
 atgcagtttc ttataaacia agcactccta tcgaa 335

<210> 135  
 <211> 506  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 135  
 tttttttttt tttaaaaatc aacagattta ttcaagtgcc tcgggcaaata aacaacaaaac 60  
 atccacaaac ataattattat tgaacttttc ctttttaaaa cttatcaaag gccttctttg 120  
 ttcctgagac tttgatcacc ttcaaaacat taaaacgaac agttttactc aaaggcctgc 180  
 attcaccgat cgtgacaata tcaccaatag agatatcacg gaaacatggc gaacagtga 240  
 cggacatgtt tttgtgacgt ttctcgtatc gacgatattt cggaacaaag tgcaataat 300  
 cagcgcgaat gacaattgtg cgtcgcattt tgttcttgat aacaacacca gtcaaaatac 360  
 ggccacgaat tgaaacattt ccagtgaag gacactttt gtcaatataa ttgccttcga 420  
 tagcctcgcg tggagtttta aatcctaacc caacattcct ccaataacga tccttatttt 480  
 tcggctnttt gccaatccct tgcgtc 506

<210> 136  
 <211> 230  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 136  
 aattcctcaa actctgccct ggctgtcctt ctcaaaaacga caccctcgct ttattatcac 60  
 ctccagtcac ctacgaaaat tctttgagag atcaagggag taattcgaca ttatggattc 120  
 ttttgttggt ttttaattgt ttatttttgc tactaatttt ccttctaatt gccgcctacc 180  
 tccgtgtcgc catttttggc tccgccccct acaaaaacca gttccgtcgt 230

<210> 137  
 <211> 216  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 137  
 acaatacac aacaacaaaa tcattgttta ccccaaatat ccaaaagtgc tcctccaact 60  
 tcaactcgctg tttgtaccaa ctctactagc tgtaattcgt ccttagctgt gccgttaatt 120  
 tctagtgaat cggaagaaag tgatgaacaa caaagacgg ggggaatggac aaatctaaca 180  
 ttattaatta ttatttctca tgattgtaaa ttgcat 216

<210> 138  
 <211> 395  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 138  
 atgcattcct gaagcaattt tgggtatgga cattgtatgc caagcaaagt ctgggtatggg 60  
 gaagacagct gtatttgtgt tggcaacact ccaacaattg actccagtgt acgggacggg 120  
 ctctgttctc gttatgtgtc aactcgcga acttgctttt caaatttcaa aggaatatga 180  
 aagatttagc aaatatatgc ccggaactaa ggtttcgggt ttctttgggt gtatgccgat 240  
 caagaaggac gaggagactt tggctaagaa cactccgcac attgttgtt gcaactccag 300  
 gcgtctgctg gcgttgggac gtacaggaca attgaagctg aaaaacatca aattcttcgt 360  
 ttttagacgaa tgtgacaaaa tgattgggga cgctg 395

<210> 139  
 <211> 591

AKK110P1

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 139

```
gaattcggcg ttgtctcggg gtccacgctc aatttcaccg aaatttttgg ggcaggcgctc 60
ctccacacca aactctgggt cattgacaac cggcactttt atctgggttc agcaaacatg 120
gactggcagt cacttactga agtcaaggaa atgggtctta tgctgttgaa ctgctcctgt 180
ttggcgtggg aactgagcaa aatatttgcg atttactggc ggattggaca gaatcacaat 240
cgcttgcccg ctgtttggcc agttttattta caatcaaaat tcaacgctca acaccaatg 300
gaaattcatt ttggacctga gccctcgcac acgtacattt cgcactcgcc tgagaagtgt 360
aacccaaagg gcagagaaca cgacctttcg gccatatgct catgcatggg aaaagccaac 420
gaatttggtc gaattgcggg aatggattat attcctgcaa caatttacct gccgaatggg 480
aacaacatat attggccatc gatcgaatgac gcgataagaa cggcagctta tcgggggtgtg 540
aaagttgacc tttggtgagt ctgtggcccc atttgaatga acgagcgatt t 591
```